

---

# ANESTHESIA & ANALGESIA

---

*Journal of the International Anesthesia Research Society, the Society of Cardiovascular Anesthesiologists,  
the Society for Pediatric Anesthesia, and the Society for Ambulatory Anesthesia*

---

Index  
Volume 87, 1998

## EDITORIAL BOARD Anesthesia & Analgesia

*Editor-in-Chief*

Ronald D. Miller, MD

*Associate Editor-in-Chief Cardiovascular Anesthesia*

Kenneth J. Tuman, MD

*Editors*

David J. Birnbach, MD, *Obstetric Anesthesia*

Thomas J. J. Blanck, MD, PhD

Daniel B. Carr, MD

Pierre Coriat, MD

Peter G. Duncan, MD, *Economics and Health Systems Research*

Thomas J. Gal, MD

Simon Gelman, MD, PhD

William J. Greeley, MD, MBA, *Pediatric Anesthesia*

Sten Lindahl, MD

Michael F. Roizen, MD

John C. Rowlingson, MD

Koh Shingu, MD

Graham Smith, BSc, MD, FRCA

Christoph Stein, MD

Peter M. Suter, MD, *Critical Care and Trauma*

Hugo Van Aken, MD, PhD

David S. Warner, MD, *Neurosurgical Anesthesia*

Denise J. Wedel, MD, *Regional Anesthesia and Pain Management*

Paul F. White, MD, PhD, *Ambulatory Anesthesia*

*Book and Multimedia Review Editor*

Norig Ellison, MD

*Statistical Editor*

Nathan L. Pace, MD

## Author Index

- Aantaa R, see Taittonen MT
- Aasboe V, Raeder JC, Groegaard B.  
Betamethasone reduces  
postoperative pain and nausea after  
ambulatory surgery, 319
- Abdi S, Lee DH, Chung JM. The anti-  
allodynic effects of amitriptyline,  
gabapentin, and lidocaine in a rat  
model of neuropathic pain, 1360
- Abe K, Shimizu T, Takashina M, Shiozaki  
H, Yoshiya I. The effects of  
propofol, isoflurane and sevoflurane  
on oxygenation and shunt fraction  
during one-lung ventilation, 1164
- Takada K, Yoshiya I. Sevoflurane and  
torsade de pointes (letter, reply),  
981  
see Komatsu H
- Abeyewardene L, see Emmons S
- Aguirre C, see Zamacona MK
- Ahmed HE, Craig WF, White PF,  
Ghoname E-SA, Hamza MA, Gajraj  
NM, Taylor SM. Percutaneous  
electrical nerve stimulation: an  
alternative to antiviral drugs for  
acute herpes zoster, 911
- Aigret D, see Godet G
- Akamatsu H, see Mikawa K
- Akamatsu S, see Ueda N
- Akatsuka M, see Doi Y
- Albani A, see Runza M
- Alberts MS, see Sniadach MS
- Aldegheri G  
see Casati A  
see Fanelli G
- Ali OS, see Joly L-M
- Allen HW, Liu SS, Ware PD, Nairn CS,  
Owens BD. Peripheral nerve blocks  
improve analgesia after total knee  
replacement surgery, 93
- Amaro AR, see Carvalho JCA
- Ambesh SP, Kaushik S. Percutaneous  
dilatational tracheostomy: the Ciaglia  
method versus the Rapi-trach  
method, 556
- Amr N, see Kasper S-M
- Andel H, see Koinig H
- Andersen J, see Callesen T
- Anderson JL, see Hill GE
- Angelides AG, see Schwartz DA
- Angelescu D, Kaplan EF. Defining levels  
of sedation in pediatric patients  
(letter), 1454
- Antognini JF, Carstens E. A simple,  
quantifiable, and accurate method  
for applying a noxious mechanical  
stimulus, 1446
- Jinks S, Buzin V, Carstens E. A method  
for differential delivery of  
intravenous drugs to the head and  
torso of the goat, 1450
- Anwar M, see Kaye AD
- Appavu SK, see Khorasani A
- Appelbaum A, see Roytblat L
- Appleyard RW. Performance-based  
credentialing, not pediatric  
subspecialization, is the issue  
(letter), 499
- Arantes FM, see Capelozzi M
- Arkoosh VA. Pain Relief in Labour. 1st ed.  
(book review), 749  
see Huffnagle SL
- Armstrong TSH, Byers GF, Johnston GM.  
Tracheal intubation in infants  
(letter), 1455
- Arndt JO, Bömer W, Krauth J, Marquardt  
B. Incidence and time course of  
cardiovascular side effects during  
spinal anesthesia after prophylactic  
administration of intravenous fluids  
or vasoconstrictors, 347
- Artru AA  
see Momota Y  
see Talmor D
- Arya VK. Understanding and defining  
minimum alveolar anesthetic  
concentration (letter), 1453
- Asai T. Use of a lighted stylet and the  
laryngeal mask for tracheal  
intubation (letter), 498
- Shingu K, Latto IP. Use of a lighted  
stylet for tracheal intubation  
through the intubating laryngeal  
mask (letter), 979
- Ashburn MA, see Rathmell JP
- Ashworth J, Smith I. Comparison of  
desflurane with isoflurane or  
propofol in spontaneously breathing  
ambulatory patients, 312
- Asokumar B, Newman LM, McCarthy RJ,  
Ivankovich AD, Tuman  
KJ. Intrathecal bupivacaine reduce  
pruritus and prolongs duration of  
fentanyl analgesia during labor: a  
prospective, randomized controlled  
trial, 1309
- Atkins RF. Allocating subspecialist  
pediatric anesthesiologists: when to  
say no (letter), 742
- Atkinson R, see Richardson J
- Augustatou CG, see Mentzelopoulos SD
- Avramov MN, Griffin JD, White PF. The  
effect of fresh gas flow and  
anesthetic technique on the ability  
to control acute hemodynamic  
responses during surgery, 666
- Ayoub CM, Ghobashy A, Koch ME,  
McGrimley L, Pascale V, Qadir S,  
Ferneini EM, Silverman DG.  
Widespread application of topical  
steroids to decrease sore throat,  
hoarseness, and cough after tracheal  
intubation, 714
- Babacan A, Öztürk E, Kaya K. Relief of  
chronic refractory hiccups with  
glossopharyngeal nerve block  
(letter), 980
- Bacconnier P, see Carry P-Y
- Badgwell JM, see Sessler DI
- Badner NH, Drader K, Freeman D, Spence  
JD. The use of intraoperative  
nitrous oxide leads to postoperative  
increases in plasma homocysteine,  
711
- Badr A, see Saha AK
- Baer GA, see Puura AIE
- Bailey MK, see Harvey SC
- Bailey S, see Bastien JL
- Bains J, see Ellis JE
- Bakhos M, see Chaney MA
- Bakker PFA, see de Vries JW
- Balea M, see Russell IAM
- Ballantyne JC, Chang Y-C. The impact of  
choice of muscle relaxant on  
postoperative recovery time (letter,  
reply), 499
- Banister RE, see Kaye AD
- Banssillon V  
see Carry P-Y  
see Preckel MP
- Barcia JA, see Gil F
- Barnett G, see Bloomfield EL
- Barnhart JM, see Katz RI
- Barry J, see Brimacombe JR
- Bartels CF, see Vanlinthout LE
- Bashour CA, see Yared J-P
- Bastanmehr H, see De Witte JL
- Bastien JL, Graves JR, Bailey S. Atypical  
presentation of amniotic fluid  
embolism, 124
- Baurain MJ, Hennart DA, Godschalx A,  
Huybrechts I, Nasrallah G,  
d'Hollander AA, Cantraine F.  
Visual evaluation of residual  
curarization in anesthetized patients  
using one hundred-hertz, five-  
second tetanic stimulation at the  
adductor pollicis muscle, 185

- Bazzell CM, see Dunn SM  
 Beccaria P, see Fanelli G  
 Bech K, see Callesen T  
 Bedell EA, Prough DS. Should induced hypertension be beneficial after traumatic brain injury? (editorial), 751  
 Beems T, see Harbers JBM  
 Beers RA, see Cucchiari G  
 Behforouz N, Dubousset AM, Jamali S, Ecoffey C. Respiratory effects of desflurane anesthesia on spontaneous ventilation in infants and children, 1052  
 Belden C, see Greenberg RS  
 Bellows WH, see Leung JM  
 Benhamou D, Thorin D, Brichant J-F, Dailland P, Milon D, Schneider M. Intrathecal clonidine and fentanyl with hyperbaric bupivacaine improves analgesia during cesarean section, 609  
 see Joly L-M  
 Benumof JL. Use of the laryngeal mask airway as an alternative to the tracheal tube during ambulatory anesthesia (letter), 977  
 Berger R, see Kozek-Langenecker SA  
 Berkowitz RJ, see Litman RS  
 Berlakovich GA, see Krenn CG  
 Berman AT, see Parmet JL  
 Berman L, see Saha AK  
 Bernard JM, Fulgencio JP, Delaunay L, Bonnet F. Clonidine does not impair redistribution hypothermia after the induction of anesthesia, 168  
 Berry AM, see Brimacombe JR  
 Berry RD, see Forde SC  
 Berti M, see Fanelli G  
 Bevan DR, see Bevan JC  
 Bevan JC, Reimer EJ, Smith MF, deV. Scheepers L, Bridge HS, Martin GR, Bevan DR. Decreased mivacurium requirements and delayed neuromuscular recovery during sevoflurane anesthesia in children and adults, 772  
 Bhardwaj A, see Harukuni I  
 Biber B, see Sundeman H  
 Bigatello L, see Viguera A  
 Birmbach DJ, see Runza M  
 Biro P, Eylich G, Rohling RG. The efficiency of CO<sub>2</sub> elimination during high-frequency jet ventilation for laryngeal microsurgery, 180  
 Bishop MJ, see Kim ES  
 Bjorksten AR, see De Witte JL  
 Blakeley KR, Klein KW, White PF, Trott S, Rohrich RJ. A total intravenous anesthetic technique for outpatient facial laser resurfacing, 827  
 Blakeman B, see Chaney MA  
 Blanck TJJ, see Xu F  
 Blanot S, see Orliaguet GA  
 Bleeker CP, see Harbers JBM  
 Bloomfield EL, Schubert A, Secic M, Barnett G, Shutway F, Ebrahim ZY. The influence of scalp infiltration with bupivacaine on hemodynamics and postoperative pain in adult patients undergoing craniotomy, 579  
 Boban N, see McCallum JB  
 Bode ET, see McNitt JD  
 Bömer W, see Arndt JO  
 Bonnet F, see Bernard JM  
 Booi LH, see Vanlinthout LE  
 Borchardt RA, LaQuaglia MP, McDowall RH, Wilson RS. Bronchial injury during lung isolation in a pediatric patient, 324  
 Borgeat A, Hasler P, Fahti M. Gynecologic laparoscopic surgery is not associated with an increase of serotonin metabolites excretion, 1104  
 Ruetsch YA, Jörg M. Convulsions induced by ropivacaine during interscalene brachial plexus block (letter), 497  
 Bornstein MH, see Kain ZN  
 Bosnjak ZJ, see McCallum JB  
 Bouaziz H, see Zetlaoui PJ  
 Boudreault D, see Couture P  
 Bourke DL, Thomas P. Mandibular nerve block in addition to cervical plexus block for carotid endarterectomy, 1034  
 Bovill JG, see Camu F  
 Bowen PJ, Windsor A, Flaishon R, Sebel PS. The interaction of etanalone and fentanyl with special reference to logistic regression analysis, 967  
 Boxer L, see Huffnagle SL  
 Boylan JF, see Lydon AM  
 Braslavsky A, see Tverskoy M  
 Bräu ME, Vogel W, Hempelmann G. Fundamental properties of local anesthetics: half-maximal blocking concentrations for tonic block of Na<sup>+</sup> and K<sup>+</sup> channels in peripheral nerve, 885  
 Brecher ME, see Monk TG  
 Brechtelsbauer H, see Orth VH  
 Bremerich DH, see Kai T  
 Brennan TJ, see Zahn PK  
 Brichant J-F, see Benhamou D  
 Bridge HS, see Bevan JC  
 Brimacombe J, Keller C. A comparison of pharyngeal mucosal pressure and airway sealing pressure with the laryngeal mask airway in anesthetized adult patients, 1379  
 Keller C, Morris R, Mecklem D. A comparison of the disposable versus the reusable laryngeal mask airway in paralyzed adult patients, 921  
 Brimacombe JC, see Brimacombe JR  
 Brimacombe JR, Brimacombe JC, Berry AM, Morris R, Mecklem D, Clarke G, Barry J, Kirk T. A comparison of the laryngeal mask airway and cuffed oropharyngeal airway in anesthetized adult patients, 147  
 Brock-Utne JG  
 see Habibi A  
 see Truelsen KS  
 Broekema AA, Veen A, Fidler V, Gielen MJM, Hennis PJ. Postoperative analgesia with intramuscular morphine at fixed rate versus epidural morphine or sufentanil and bupivacaine in patients undergoing major abdominal surgery, 1346  
 Bromberg M, see Swenson JD  
 Bronsther O, see Lustik SJ  
 Brook M, see Russell IAM  
 Broomé M, see Sundeman H  
 Brown DL, see Wilson JL  
 Brown RS, see Mongan PD  
 Bruda NL, Hofmann SC. Controlled continuous spinal drainage using a flow-regulated pump, 1306  
 Brüssel T, see Knichwitz G  
 Bryson GL. The efficacy of acute normovolemic hemodilution (letter, reply), 498  
 Buerkle H  
 see Marcus MAE  
 see Tsai PS  
 Buisson C, see Orliaguet GA  
 Bund M, see Walz R  
 Bushell E, see Habibi A  
 Butterworth JF, see Ririe DG  
 Buzelin J-M, see Malinovsky J-M  
 Buzello W, see Kasper S-M  
 Buzin V, see Antognini JF  
 Byers GF, see Armstrong TSH  
 Cahalan MK, see Russell IAM  
 Cahill DR, see Wilson JL  
 Callens K, see Vanlinthout LE  
 Callesen T, Bech K, Thorup J, Andersen J, Nielsen R, Roikjær O, Kehlet H. Cryoanalgesia: effect on postherniorrhaphy pain, 896  
 Calvo R, see Zamacona MK



- Camann W, see Oh J  
Camann WR, see Levin A  
Campbell CE, see Griffin PR  
Camu F, Van Aken H, Bovill JG.  
Postoperative analgesic effects of  
three demand-dose sizes of fentanyl  
administered by patient-controlled  
analgesia, 890  
Canessa R, see Lema G  
Cantraine F, see Baurain MJ  
Capan LM, Sutin K, Ibarra PF, Wahlander  
S. Succinylcholine cannot relieve an  
airway obstruction caused by  
pharyngeal and laryngeal edema  
(letter, reply), 229  
Capdeville M, Hall D, Koch CG. Practical  
use of a bronchial blocker in  
combination with a double-lumen  
endotracheal tube, 1239  
Capehart BP, see Pisetsky MA  
Capelozzi M, Arantes FM, Paiva PSO,  
Capelozzi VL, Martins MA. Spinal  
anesthesia increases pulmonary  
responsiveness to methacholine in  
guinea pigs, 874  
Capelozzi VL, see Capelozzi M  
Cappelleri G, see Casati A  
Cappelli EL, see Carvalho JCA  
Caramico LA, see Kain ZN  
Cardoso MMSC, see Carvalho JCA  
Carli PA, see Orliaguet GA  
Carmona JA, see De La Cruz JP  
Carpenter R, see Gerancher JC  
Carroll MT, see Lowry DW  
Carry P-Y, Gallet D, François Y, Perdrix  
J-P, Sayag A, Gilly F, Eberhard A,  
Bansillon V, Baconnier P.  
Respiratory mechanics during  
laparoscopic cholecystectomy: the  
effects of the abdominal wall lift,  
1393  
Carstens E, see Antognini JF  
Carvalho JCA, Amaro AR, Cappelli EL,  
Cardoso MMSC, Prado AA.  
Combination of diclofenac and  
intrathecal morphine for cesarean  
delivery (letter, reply), 1453  
Casaletti E, see Casati A  
Casati A, Fanelli G, Cappelleri G,  
Aldegheri G, Leoni A, Casaletti E,  
Torri G. Effects of spinal needle  
type on lateral distribution of 0.5%  
hyperbaric bupivacaine, 355  
see Fanelli G  
Castellani WJ, see Sprung J  
Castro AD, see Gaiser RR  
Castro B, see Mazurek AJ  
Cavill G, see Mahajan RP  
Chakrabarti MK, see Ma D  
Chamberlain SK, see Shaw ADS  
Champsaur G, see Vedrinne JM  
Chan MTV, Chui PT, Ho WS, King WWK.  
Single-dose tropisetron for  
preventing postoperative nausea  
and vomiting after breast surgery,  
931  
Chaney MA, Nikolov MP, Blakeman B,  
Bakhos M, Slogoff S. Pulmonary  
effects of methylprednisolone in  
patients undergoing coronary artery  
bypass grafting and early tracheal  
extubation, 27  
Chang Y-C, see Ballantyne JC  
Charchafieh J, Cottrell JE, Kass IS. The  
effect of fentanyl on  
electrophysiologic recovery of CA 1  
pyramidal cells from anoxia in the  
rat hippocampal slice, 68  
Chari P  
see Puri GD  
see Thiagarajan A  
Charron B, see Orliaguet GA  
Chase GA, see Hannallah MS  
Chavez R, see Wittels B  
Chelliah RY, see Chow MYH  
Chen EH, see Tobin MJ  
Chen L, Tang J, White PF, Sloninsky A,  
Wender RH, Naruse R, Kariger R.  
The effect of location of  
transcutaneous electrical nerve  
stimulation on postoperative opioid  
analgesic requirement: acupoint  
versus nonacupoint stimulation,  
1129  
Cheng EY, see Finck M  
Cheng C-H, see Wong C-S  
Chhibber AK, see Lustik SJ  
Chin Y-J, see Chung C-J  
Chodak GW, see Worwag E  
Choe H. New epidural drugs: *primum non  
nocere* (letter, reply), 1212  
Chow MYH, Liam BL, Lew TWK, Chelliah  
RY, Ong BC. Predicting the size of a  
double-lumen endobronchial tube  
based on tracheal diameter, 158  
Chow TKF, Penberthy AJ, Goodchild CS.  
Ketamine as an adjunct to morphine  
in postthoracotomy analgesia: an  
unintended N-of-1 study, 1372  
Chui PT, see Chan MTV  
Chung C-J, Kim J-S, Park H-S, Chin Y-J.  
The efficacy of intrathecal  
neostigmine, intrathecal morphine,  
and their combination for post-  
cesarean section analgesia, 341  
Chung F, see Jin F  
Chung JM, see Abdi S  
Clarke G, see Brimacombe JR  
Cobby TF, see Crighton IM  
Cohen AM, see Pivalizza EG  
Cohen DE, see Rose JB  
Cohen I, see Ezri T  
Cohen MM, see Fung D  
Cohen N, see Miller RD  
Collins HV, see Richardson MG  
Collins L, see Parmet JL  
Colonna-Romano P, see Parmet JL  
Connelly NR  
see Dunn SM  
see Reuben SS  
see Schwartz DA  
Cook-Sather SD, Nicolson SC, Schreiner  
MS. Tracheal intubation in infants  
(letter, reply), 1456  
Coraim F, see Kozek-Langenecker SA  
Coriat P  
see Godet G  
see Lejay M  
Cosgrove DM III, see Yared J-P  
Coté CJ  
see Mazurek AJ  
see Tobin MJ  
Cottrell JE, see Charchafieh J  
Cousins MJ, see Grace RF  
Couture P, Boudreault D. Epidural  
analgesia and intravenous patient-  
controlled analgesia result in similar  
rates of myocardial ischemia after  
aortic surgery (letter, reply), 745  
Cozian A, see Malinovsky J-M  
Craig WF, see Ahmed HE  
Crain B, see Harukuni I  
Crighton IM, Martin PH, Hobbs GJ, Cobby  
TF, Fletcher AJP, Stewart PD. A  
comparison of the effects of  
intravenous tramadol, codeine, and  
morphine on gastric emptying in  
human volunteers, 445  
Crocì M, see Pelosi P  
Crystal GJ, Zhou X, Gurevicius J, Salem  
MR. Influence of nitric oxide on  
vascular, metabolic, and contractile  
responses to dobutamine in *in situ*  
canine hearts, 994  
Csavoy AN, see Segal S  
Cucchiari G, Beers RA. Recrudescence of  
opioid effects after intraoperative  
remifentanyl infusion complicated  
by infiltration of the intravenous  
catheter (letter), 1458  
Curry R, see Eisenach JC  
Curtis A, see Vedrinne JM  
Czeslick E, see Doenicke AW  
Daghfous M, see Godet G  
Dailland P, see Benhamou D  
Daley MD, Norman PH. Torsade de  
pointes ventricular tachycardia

- during, but not necessarily caused by, sevoflurane anesthesia (letter), 981
- Damask MC, see Pagel PS
- Dandona P, see Nader-Djalal N
- D'Angelo R  
see Gage JC  
see Gerancher JC  
see Tang J
- Datta S  
see Levin A  
see Segal S
- Davidson BA, see Nader-Djalal N
- Davis RF, see Pagel PS
- Dayan SS, see Katz RI
- De Amici D, see Ramajoli F
- de Blok S, see van Ee R
- Deem S. Sevoflurane and torsade de pointes (letter), 981
- Deepalakshmi K, see Kundra P
- De Kock M  
see Von Montigny S  
see Quintin L
- De La Cruz JP, Sedeño G, Carmona JA, Sánchez de la Cuesta F. The *in vitro* effects of propofol on tissular oxidative stress in the rat, 1141
- de Leon-Casasola OA, Lema MJ. Epidural analgesia and intravenous patient-controlled analgesia result in similar rates of myocardial ischemia after aortic surgery (letter), 745
- Delaunay L, see Bernard JM
- Delbrouck D, see Godet G
- de Vries JW, Bakker PFA, Visser GH, Diephuis JC, van Huffelen AC. Changes in cerebral oxygen uptake and cerebral electrical activity during defibrillation threshold testing, 16
- De Witte JL, Kim J-S, Sessler DI, Bastanmehr H, Bjorksten AR. Tramadol reduces the sweating, vasoconstriction, and shivering thresholds, 173
- DeLong ER  
see Klein SM  
see Lubarsky DA
- Depper JA, see Palmon SC
- D'Ercole FJ  
see Klein SM  
see Steele SM
- Deroy R, Ghoris M. The Combitude elective anesthetic airway management in a patient with cervical spine fracture, 1441
- deV. Scheepers L, see Bevan JC
- Dexter F  
see Ludington E
- see Sarraf-Yazdi S
- Deyaert M, see Singelyn FJ
- d'Hollander AA, see Baurain MJ
- Dickstein ML, see Heerdt PM
- Diephuis JC, see de Vries JW
- Dodd-o JM. Safety of  $\alpha_2$ -agonists? (letter, reply), 496
- Dodson BA. Clinical Neuroanesthesia. 2nd ed. (book review), 983
- Doenicke AW, Czeslick E, Moss J, Hoerneck R. Onset time, endotracheal intubating conditions, and plasma histamine after cisatracurium and vecuronium administration, 434
- Dohi K, see Kikumoto K
- Dohi S, see Ueda N
- Doi Y, Uda R, Akatsuka M, Tanaka Y, Kishida H, Mori H. Damaged Univent tubes, 732
- Done ML, Lee A. The use of a video to convey preanesthetic information to patients undergoing ambulatory surgery, 531
- Donnelly M, see Duncan M
- Donzelli J, see Pappas ALS
- Dorje P, see LaGorio J
- Douglas MJ, see Pavy TJG
- Drader K, see Badner NH
- Drage M. Caution in the use of lidocaine infusion in the surgical patient (letter), 1212
- Driver RP Jr. Prefilled syringes should be available in obstetric units and operating rooms (letter, reply), 1457
- Drop LJ, see Viguera A
- Dubey PK, see Sinha PK
- Dubousset AM, see Behforouz N
- Dubreuil C, see Preckel MP
- Duclaux R, see Preckel MP
- Duke JH Jr, see Pivalizza EG
- Dull RO, Peterfreund RA. Variations in the composition of spinal anesthetic solutions: the effects of drug addition order and preparation methods, 1326
- Duncan M, Donnelly M. A comparison of the antiemetic effect of ondansetron and droperidol on patients undergoing gynecologic laparoscopy (letter), 1213
- Dunn SM, Connelly NR, Steinberg RB, Lewis TJ, Bazzell CM, Klatt JL, Parker RK. Intrathecal sufentanil versus epidural lidocaine with epinephrine and sufentanil for early labor analgesia, 331  
see Schwartz DA
- Duong DH, see Joshi S
- Durieux ME, see Kohrs R
- Duvdenani A, see Roytblat L
- Ebel C, see Holst D
- Eberhard A, see Carry P-Y
- Ebrahim ZY, see Bloomfield EL
- Eckard JB, see Johnson RF
- Eckenhoff RG. A noble approach to mechanisms (letters), 239
- Ecoffey C, see Behforouz N
- Egawa H, see Takiguchi T
- Eger EI II, Weiskopf RB. No evidence of sevoflurane-induced renal injury in volunteers (letter, reply), 230  
see Gong D  
see Koblin DD  
see Sonner JM  
see Trudell JR
- Ehman RL, see Wilson JL
- Eichelberger JP, see Lustik SJ
- Eisenach JC, Hood DD, Curry R. Intrathecal, but not intravenous, clonidine reduces experimental thermal or capsaicin-induced pain and hyperalgesia in normal volunteers, 591  
James FM III, Gordh T Jr, Yaksh TL. New epidural drugs: *primum non nocere* (letter), 1211
- Eledjam J-J, see Ripart J
- Eliasziw M, see Watts ADJ
- Ellis JE, Pedlow S, Bains J. Premedication with clonidine does not attenuate suppression of certain lymphocyte subsets after surgery, 1426
- Ellison N. Perioperative Transfusion Medicine (book review), 235
- Emmons S, Abeyewardene L, Ramakrishnan U. Case of frozen thiopental (letter), 748
- Engelhard K, see Lu H
- Erasmí H, see Kasper S-M
- Erb JM. A less difficult method to protect the tracheal cuff of a double-lumen tube (letter), 1217
- Erb T, Marsch SC, Hampl KF, Frei KJ. Fiberoptic intubation and children (letter, reply), 495
- Eriator II. Naloxone: acute opioid withdrawal syndrome or side effects? (letter), 1214
- Evers AS, see Waddle JP
- Eyrich G, see Biro P
- Ezri T, Cohen I, Geva D, Szmuk P. Pharyngoscopic views (letter), 748
- Fahti M, see Borgeat A
- Fanelli G, Casati A, Beccaria P, Aldegheri G, Berti M, Tarantino F, Torri G. A double-blind comparison of

- ropivacaine, bupivacaine, and mepivacaine during sciatic and femoral nerve blockade, 597  
see Casati A
- Fang Z  
see Gong D  
see Koblin DD
- Faure EAM, see Wittels B
- Faust RJ. Acute normovolemic hemodilution versus predonation: what is the actual cost? (letter), 233
- Feng CJ, see Kaye AD
- Ferber-Viart C, see Preckel MP
- Ferman R, see Tverskoy M
- Ferneini EM, see Ayoub CM
- Ferouz F, Norris MC. Mechanism of analgesic action of intrathecal sufentanil (letter), 1211  
see Huffnagle SL
- Ferrero-Conover D, see Ross AK
- Fidler V, see Broekema AA
- Fieber T, see Konrad Ch
- Finck M, Cheng EY. Acute pseudotongue enlargement after general anesthesia, 1443
- Fineman JR, see Russell IAM
- Finn HA, see Renz CL
- Finsterer U, see Orth VH
- Fiset P, see Tran T-V
- Flaishon R  
see Bowen PJ  
see Iselin-Chaves IA
- Fleisher LA, Mantha S, Roizen MF. Medical technology assessment: an overview, 1271
- Fletcher AJP, see Crighton IM
- Fogel ST, Shyken JM, Leighton BL, Mormol JS, Smeltzer JS. Epidural labor analgesia and the incidence of cesarean delivery for dystocia, 119
- Forde SC, Berry RD. Prefilled syringes should be available in obstetric units and operating rooms (letter), 1457
- Fournier R, Gamulin Z, Van Gessel E. Respiratory depression after 5 µg of intrathecal sufentanil, 1377
- Fox L, see Schug SA
- Framstad M, see Nguyen AC
- Franck M, see Vedrinne JM
- François Y, see Carry P-Y
- Frank DU, Horstman DJ, Rich GF. The effect of prolonged inhaled nitric oxide on pulmonary vasoconstriction in rats, 1285
- Frass M  
see Hofbauer R  
see Krafft P
- Freeman D, see Badner NH
- Frei FJ, see Erb T
- Friemann S, see Müller M
- Froelich EG, see Greilich PE
- Fu JL, see Pagel PS
- Fujii Y, Saitoh Y, Tanaka H, Toyooka H. Prophylactic oral antiemetics for preventing postoperative nausea and vomiting: granisetron versus domperidone, 1404
- Takahashi S, Toyooka H. The effects of milrinone and its mechanism in the fatigued diaphragm in dogs, 1077
- Toyooka H, Tanaka H. A granisetron-droperidol combination prevents postoperative vomiting in children, 761
- Fujise K, Shingu K, Matsumoto S, Nagata A, Mikami O, Matsuda T. The effects of the lateral position on cardiopulmonary function during laparoscopic urological surgery, 925
- Fukuda K, Kato S, Shoda T, Morikawa H, Mima H, Mori K. Partial agonistic activity of naloxone on the opioid receptors expressed from complementary deoxyribonucleic acids in Chinese hamster ovary cells, 450
- Fulgencio JP, see Bernard JM
- Fung D, Cohen MM. Measuring patient satisfaction with anesthesia care: a review of current methodology, 1089
- Furuya H  
see Kawaguchi M  
see Kikumoto K  
see Sha K
- Gage JC, D'Angelo R. Mechanism of analgesic action of intrathecal sufentanil (letter, reply), 1211
- Gaiser RR, Castro AD. The level of anesthesia resident training does not affect the risk of dental injury, 255
- Gajraj NM, see Ahmed HE
- Gallet D, see Carry P-Y
- Galley HF, see Le Cras AE
- Gambling DR, see Pavy TJG
- Gamulin Z  
see Fournier R  
see Niquille M
- Gan TJ, Rogers S. Informed use of buccal oximetry (letter, reply), 495  
see Iselin-Chaves IA
- Gandhi CD, see Heerdt PM
- Ganeshram T, see Jin F
- García E, see Zamacona MK
- García-Aguado R, see Gil F
- Garnerin P, see Tramèr MR
- Garry B, Woo P, Perrault DF Jr, Shapshay SM, Wurm WH. Jet ventilation in upper airway obstruction: description and model lung testing of a new jetting device, 915
- Gattinoni L, see Pelosi P
- Gelb AW, see Watts ADJ
- Genevro JL, see Kain ZN
- Gerancher JC, D'Angelo R, Carpenter R. Caudal epidural blood patch for the treatment of postdural puncture headache, 394
- Gerber H, see Konrad Ch
- Gerdas E, see Smith CE
- Geva D, see Ezri T
- Gharagozloo F, see Hannallah MS
- Ghignone M, see Quintin L
- Ghobashy A, see Ayoub CM
- Ghoname E-SA, see Ahmed HE
- Ghoneim MM. Transient global amnesia: a cause for postanesthetic memory disorder (letter), 980
- Ghoris M, see Deroy R
- Gibson CS  
see Reuben SS  
see Schwartz DA
- Gielen MJM, see Broekema AA
- Giffard RG, see Habibi A
- Gil F, García-Aguado R, Barcia JA, Guijarro E, Hostalet F, Tommasi-Rosso M, Grau F. The effect of fibrin glue patch in an *in vitro* model of postdural puncture leakage, 1125
- Gilly F, see Carry P-Y
- Ginsberg B, see Iselin-Chaves IA
- Glaser C, see Kozek-Langenecker SA
- Glass PSA, see Iselin-Chaves IA
- Gleason D, see Steele SM
- Gleason DH, see Klein SM
- Godet G, Gossens S, Prayssac P, Daghfous M, Delbrouck D, Aigret D, Coriat P. Infusion of propofol, sufentanil, or midazolam for sedation after aortic surgery: comparison of oxygen consumption and hemodynamic stability, 272
- Godschalx A, see Baurain MJ
- Gogarten W, see Marcus MAE
- Gomes MN, see Hannallah MS
- Gong D, Fang Z, Ionescu P, Laster MJ, Terrell RC, Eger EI II. Rat strain minimally influences anesthetic and convulsant requirements of inhaled compounds in rats, 963  
see Koblin DD
- Gonzales J, see Parmet JL
- Goodchild CS, see Chow TKF
- Goodman NW. Neither obsession nor

- distraction: words must be chosen well (letter), 743
- Goodnough LT, see Monk TG
- Gordh T Jr, see Eisenach JC
- Gordon J, see Liguori GA
- Gordon MA, see Liguori GA
- Gossens S, see Godet G
- Goto T, see Nakata Y
- Gouverneur JM, see Singelyn FJ
- Goverdhan PD, see Thiagarajan A
- Goyagi T, Tanaka M, Nishikawa T. Oral clonidine premedication enhances the pressor response to ephedrine during spinal anesthesia, 1336
- Grace RF, Power I, Umedaly H, Zammit A, Mersiades M, Cousins MJ, Mather LE. Preoperative dextromethorphan reduces intraoperative but not postoperative morphine requirements after laparotomy, 1135
- Gralow I. The Management of Pain (book review), 237
- Granry JC, see Preckel MP
- Grau F, see Gil F
- Graves JR, see Bastien JL
- Greeley WJ. Clinical Pediatric Anesthesia (book review), 236
- Greenberg L, see Roytblat L
- Greenberg RS, Zahurak M, Belden C, Tunkel DE. Assessment of oropharyngeal distance in children using magnetic resonance imaging, 1048
- Greengrass RA  
see Klein SM  
see Steele SM
- Greilich PE, Randle DW, Froelich EG, Yee LL. Massive intraoperative pulmonary embolism coincident with the administration of succinylcholine, 491
- Griffin JD, see Avramov MN
- Griffin PR, Mitchell MR, Viswanathan S, Riopelle JM, Campbell CE, Karpan DJ. Use of plastic rod/sleeve combination to facilitate double- to single-lumen tracheal tube exchange in patients with difficult glottic visualization (letter), 744
- Grocott HP. Atlas of Transesophageal Echocardiography (book review), 749
- Groegaard B, see Aasboe V
- Grouidine S. Informed use of buccal oximetry (letter), 494
- Lumb PD. Caution in the use of lidocaine infusion in the surgical patient (letter, reply), 1213
- Lumb PD. Other ways to stimulate postoperative bowel function (letter, reply), 500
- Gruber E, see Tschernko EM
- Grüne F, see Kasper S-M
- Guijarro E, see Gil F
- Guo TZ, see Poree LR
- Gupta SK, see Hooda S
- Gurevicius J, see Crystal GJ
- Gurman GM, see Roytblat L
- Habibi A, Bushell E, Jaffe RA, Giffard RG, Brock-Utne JG. Two tips for users of Bullard intubating laryngoscope, 1206
- Hacein-Bey L, see Joshi S
- Hachenberg T, see Sielenkämper AW
- Haddadin AS, see Shelley KH
- Hagen JF, see Smith CE
- Häggmark S, see Sundeman H
- Hahn MB. Management of Acute Pain. Vol. 10, No. 1 of Problems in Anesthesia (book review), 984
- Haider W, see Tschernko EM
- Haiek M, see Runza M
- Hall D, see Capdeville M
- Hall SC  
see Rockoff MA  
see Tobin MJ
- Haller M, see Orth VH
- Halsey MJ, see Koblin DD
- Hamaya Y, see Ueda N
- Hammon JW Jr, see Ririe DG
- Hampl KF, see Erb T
- Hamza MA, see Ahmed HE
- Hanaoka K, see Nishiyama T
- Haney M, see Sundeman H
- Hanley FL, see Russell IAM
- Hann S, see Mazurek AJ
- Hannallah MS, Gharagozloo F, Gomes MN, Chase GA. A comparison of the reliability of two techniques of left double-lumen tube bronchial cuff inflation in producing water-tight seal of the left mainstem bronchus, 1027
- Hanouz J-L, see Lejay M
- Hanson EW, Lowson SM. Atrial fibrillation and thromboembolism, 217
- Hao J-X, see von Heijne M
- Hara K, Saito Y, Kirihaara Y, Sakura S, Kosaka Y. Antinociceptive effects of intrathecal L-type calcium channel blockers on visceral and somatic stimuli in the rat, 382
- Harbers JBM, Beems T, Hoen MB, Bleeker CP. A case of temporary facial nerve palsy after regional anesthesia of the scalp, 1375
- Hartmann T, see Preis CA
- Hartung J. Emesis and nitrous oxide: a correction (letter), 1215
- Harukuni I, Bhardwaj A, Traystman RJ, Crain B, London ED, Kirsch JR. Neuroprotection from focal ischemia by 4-phenyl-1-(4-phenylbutyl) piperidine (PPBP) is dependent on treatment duration in rats, 1299
- Harvey SC, Roland P, Bailey MK, Tomlin MK, Williams A. A randomized, double-blind comparison of rocuronium, *d*-tubocurarine, and "mini-dose" succinylcholine for preventing succinylcholine-induced muscle fasciculations, 719
- Hasler P, see Borgeat A
- Hastings RH, see Ng M
- Hausman R, see Holst D
- Hayashi Y, see Kikimoto K
- Heerdt PM. Intrapulmonary shunt fraction and prostaglandin E<sub>1</sub> (letter), 233
- Gandhi CD, Dickstein ML. Disparity of isoflurane effects on left and right ventricular afterload and hydraulic power generation in swine, 511
- Heidegger T, Kreienbühl G. Asystole under hypotensive epidural anesthesia (letter, reply), 982
- Hein HAT, see Ozelsel TJP
- Hempelmann G  
see Bräu ME  
see Müller M
- Hemrika DJ, see van Ee R
- Hennart DA, see Baurain MJ
- Hennis PJ, see Broekema AA
- Hersch D, see Katz RI
- Herschman Z. Other ways to stimulate postoperative bowel function (letter), 500
- Hertz R, see Saff GN
- Hettrick DA, Kersten JR, Pagel PS, Warltier DC. Interpretation of nonpulsatile arterial pressure-flow relations during cardiopulmonary bypass, 743
- Hibbard J, see Wittels B
- Higgins M, see Saha AK
- Hill GE, Anderson JL, Lyden ER. Ketamine inhibits the proinflammatory cytokine-induced reduction of cardiac intracellular cAMP accumulation, 1015
- Hines M, see Ririe DG
- Hirakawa M, see Yokoyama M
- Hirano T, see Kawana S
- Hirota K, see Sato T
- Hirsch BZ, see Schwartz DA
- Ho G, see Katz RI

- Ho KM, see Ku PKM  
 Ho S-T, see Wang J-J  
 Ho WS, see Chan MTV  
 Hobbs GJ, see Crighton IM  
 Hoen MB, see Harbers JBM  
 Hoerneck R, see Doenicke AW  
 Hofbauer R, Moser D, Salfinger H, Frass M, Kapiotis S. Sufentanil inhibits migration of human leukocytes through human endothelial cell monolayers, 1181  
 Hoffman-Hogg L, see Yared J-P  
 Hofmann SC, see Bruda NL  
 Hofstadter MB, see Kain ZN  
 Hogan Q, see McCallum JB  
 Holcomb GW, see Pretorius M  
 Holm BA, see Nader-Djalal N  
 Holst D, Möllmann M, Ebel C, Hausman R, Wendt M. *In vitro* investigation of cerebrospinal fluid leakage after dural puncture with various spinal needles, 1331  
 Hood DD, see Eisenach JC  
 Hooda S, Gupta SK. Acute transient sialadenopathy associated with laryngeal mask airway, 1438  
 Hörauf K, see Koinig H  
 Horn BJ, see Tobin MJ  
 Horrow JC, see Parmet JL  
 Horstman DJ, see Frank DU  
 Hosotsubo K, see Shimaoka M  
 Hostalet F, see Gil F  
 Hotaling AJ, see Pappas ALS  
 Hovorka J, Korttila K, Nelskylä K.  
     Reversal of neuromuscular blockade (letter, reply), 494  
 Howell S, see Iselin-Chaves IA  
 Howie MB, Kramer MG, McSweeney TD.  
     The effects of verapamil and diltiazem pretreatment on potassium release in dogs after the administration of succinylcholine, 691  
     see Pagel PS  
 Huang LT, see Tsai PS  
 Hudson RJ, see Thomson IR  
 Huffnagle HJ, see Huffnagle SL  
 Huffnagle SL, Norris MC, Arkoosh VA, Huffnagle HJ, Ferouz F, Boxer L, Leighton BL. The influence of epidural needle bevel orientation on spread of sensory blockade in the laboring parturient, 326  
 Hung OR. Use of a lighted stylet and the laryngeal mask for tracheal intubation (letter, reply), 498  
 Hunt TE, Wu W-H, Zbuzek VK. The effects of serotonin biosynthesis inhibition on nicotine and nifedipine-induced analgesia in rats, 1109  
 Hutchinson DT, see Swenson JD  
 Huybrechts I, see Baurain MJ  
 Hynynen M, see Ranta SO-V  
 Ibarra PF, see Capan LM  
 Ichinose F, see Nakata Y  
 Ikuta Y, see Shimoda O  
 Ilett KF, see Paech M  
 Imamura M, see Iwashita H  
 Inada T, see Nagata A  
 Inomata S, Suwa T, Toyooka H, Suto Y.  
     End-tidal sevoflurane concentration for tracheal extubation and skin incision in children, 1263  
 Inoue S, see Kikumoto K  
 Insler SR, see Yared J-P  
 Ionescu P  
     see Gong D  
     see Koblin DD  
 Iseki M, Okuno S, Tanabe Y, Mitsuhashi T, Miyazaki T. Methicillin-resistant *Staphylococcus aureus* sepsis resulting from infection in paravertebral muscle after continuous epidural infusion for pain control in a patient with herpes zoster, 116  
 Iselin-Chaves IA, Flaishon R, Sebel PS, Howell S, Gan TJ, Sigl J, Ginsberg B, Glass PSA. The effect of the interaction of propofol and alfentanil on recall, loss of consciousness, and the bispectral index, 949  
 Ismail M, see Wittels B  
 Isohanni MH, see Palkama VJ  
 Ivankovich AD, see Asokumar B  
 Iwasaka T, see Sha K  
 Iwasaki H, see Kawana S  
 Iwashita H, Matsukawa T, Ozaki M, Sessler DI, Imamura M, Kumazawa T. Hypoxemia decreases the shivering threshold in rabbits anesthetized with 0.2 minimum alveolar anesthetic concentration isoflurane, 1408  
 Jackson RW, see Ozelsel TJP  
 Jaffe RA, see Habibi A  
 Jamali S, see Behforouz N  
 James FM III, see Eisenach JC  
 Jamnicki M, Zollinger A, Seifert B, Popovic D, Pasch T, Spahn DR.  
     Compromised blood coagulation: an *in vitro* comparison of hydroxyethyl starch 130/0.4 and hydroxyethyl starch 200/0.5 using thrombelastography, 989  
 Jandrasits O, see Tschernko EM  
 Janfaza DR, Michna E, Pisini JV, Ross EL.  
     Bedside implantation of a trial spinal cord stimulator for intractable anginal pain, 1242  
 Jarreau M-M, see Orliaguet GA  
 Javorski JJ, see Pappas ALS  
 Jensen NF, Weiler JM. C1 esterase inhibitor deficiency, airway compromise, and anesthesia, 480  
 Jin F, Norris A, Chung F, Ganeshram T. Should adult patients drink fluids before discharge from ambulatory surgery?, 306  
 Jinks S, see Antognini JF  
 Johansen JW, Schneider G, Windsor AM, Sebel PS. Esmolol potentiates reduction of minimum alveolar isoflurane concentration by alfentanil, 671  
 Johansson G, see Sundeman H  
 Johnson KJ, see Nader-Djalal N  
 Johnson RF, Lobato EB, Eckard JB.  
     Perioperative management of a patient with latex allergy undergoing heart transplantation, 304  
 Johnston GM, see Armstrong TSH  
 Jöhr M, see Konrad Ch  
 Joly L-M, Spaulding C, Monchi M, Ali OS, Weber S, Benhamou D. Topical lidocaine-prilocaine cream (EMLA) versus local infiltration anesthesia for radial artery cannulation, 403  
 Jones J, see Richardson J  
 Jones JG, see Navapurkar VU  
 Jones KA, see Kai T  
 Jörg M, see Borgeat A  
 Joris D, see Singelyn FJ  
 Joshi GP, White PF. Use of the laryngeal mask airway as an alternative to the tracheal tube during ambulatory anesthesia (letter, reply), 978  
 Joshi S, Young WL, Pile-Spellman J, Duong DH, Vang MC, Haccin-Bey L, Lee HT, Ostapovich N. The feasibility of intracarotid adenosine for the manipulation of human cerebrovascular resistance, 1291  
 Juneja HS, see Pivalizza EG  
 Jurna I, see Niv D  
 Kadowitz PJ, see Kaye AD  
 Kahn RL, see Liguori GA  
 Kai T, Bremerich DH, Jones KA, Warner DO. Drug-specific effects of volatile anesthetics on Ca<sup>2+</sup> sensitization in airway smooth muscle, 425  
 Kain ZN, Caramico LA, Mayes LC, Genevro JL, Bornstein MH, Hofstadter MB. Preoperative



- preparation programs in children: a comparative examination, 1249
- see Kakiuchi Y  
see Miyabe M  
see Rosenblatt WH
- Kampine JP, see McCallum JB
- Kanaya N, Kawana S, Tsuchida H, Miyamoto A, Ohshika H, Namiki A. Comparative myocardial depression of sevoflurane, isoflurane, and halothane in cultured neonatal rat ventricular myocytes, 1041
- Kanbak M. Heparin resistance, antithrombin III treatment, and activated clotting time values (letter), 1215
- Phenylephrine, dopamine, mannitol, and renal protection during cardiopulmonary bypass (letter), 1458
- Kandel L, see Sonner JM
- Kaneko M, see Saito Y
- Kangawa K, see Kikumoto K
- Kanto JH, see Taittonen MT
- Kaplotis S, see Hofbauer R
- Kaplan RF, see Anghelescu D
- Kariger R, see Chen L
- Karl L, see Wittels B
- Karpan DJ, see Griffin PR
- Kasper S-M, Grüne F, Walter M, Amr N, Erasmi H, Buzello W. The effects of increased doses of bovine hemoglobin on hemodynamics and oxygen transport in patients undergoing preoperative hemodilution for elective abdominal aortic surgery, 284
- Kass IS, see Charchafliet J
- Katchko L, see Talmor D
- Kato S, see Fukuda K
- Katz RI, Barnhart JM, Ho G, Hersch D, Dayan SS, Keehn L. A survey on the intended purposes and perceived utility of preoperative cardiology consultations, 830
- Kaushik S  
see Ambesh SP  
see Sinha PK
- Kawaguchi M, Furuya H. Preferential effect of nitroglycerin on large microvessels (letter, reply), 747
- see Sha K
- Kawana S, Matsuno A, Nakabayashi K, Yamamoto S, Iwasaki H, Watanabe H, Namiki A, Hirano T. A high, large epiglottis disturbs proper positioning of the laryngeal mask and cuffed oropharyngeal airway, 489
- see Kanaya N
- Kaya K, see Babacan A
- Kaye AD, Banister RE, Anwar M, Feng CJ, Kadowitz PJ, Nossaman BD. Pulmonary vasodilation by ketamine is mediated in part by L-type calcium channels, 956
- Kaynar AM, Martinez-Ruiz R. Interluminal connection in a failed central venous line (letter), 744
- Keehn L, see Katz RI
- Kehlet H, see Callesen T
- Keller C, see Brimacombe J
- Kern FH. Echocardiography in Pediatric Heart Disease. 2nd ed. (book review), 1218
- Kersten JR, see Hettrick DA
- Kettner SC, see Kozek-Langenecker SA
- Keyes H, see Pavlin DJ
- Khorasani A, Appavu SK, Saatee S. Succinylcholine cannot relieve an airway obstruction caused by pharyngeal and laryngeal edema (letter), 229
- Kick O. The efficacy of acute normovolemic hemodilution (letter), 497
- Kihara S, see Miyabe M
- Kikumoto K, Kubo A, Hayashi Y, Minamino N, Inoue S, Dohi K, Kitamura K, Kangawa K, Matsuo H, Furuya H. Increased plasma concentration of adrenomedullin in patients with subarachnoid hemorrhage, 859
- Kim ES, Bishop MJ. Cough during emergence from isoflurane anesthesia, 1170
- Kim JL, see Mazurek AJ
- Kim J-S  
see Chung C-J  
see De Witte JL
- Kimura H, see Nagata A
- King WWK, see Chan MTV
- Kingery WS, see Poree LR
- Kirihara Y  
see Hara K  
see Saito Y
- Kirk T, see Brimacombe JR
- Kirsch JR  
see Harukuni I  
see Palmon SC
- Kirvelä OA, see Taittonen MT
- Kishida H, see Doi Y
- Kissin I, see Tverskoy M
- Kitaguchi K, see Sha K
- Kitajima T, see Takiguchi T
- Kitamura K, see Kikumoto K
- Kiyono H, see Shimaoka M
- Kizilkaya M, see Kucuk N
- Klatt JL, see Dunn SM
- Klatt M, see Reuben SS
- Klein KW, see Blakeley KR
- Klein SM, Greengrass RA, Steele SM, D'Ercole FJ, Speer KP, Gleason DH, DeLong ER, Warner DS. A comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.75% ropivacaine for interscalene brachial plexus block, 1316
- see Steele SM
- Klepckio H, see Tschernko EM
- Klimscha W, see Tschernko EM
- Knichwitz G, Van Aken H, Brüssel T. Gastrointestinal monitoring using measurement of intramucosal  $P_{CO_2}$ , 134
- Knight PR III, see Nader-Djalal N
- Kobayashi T, see Tsubokawa T
- Koblin DD, Fang Z, Eger EI II, Laster MJ, Gong D, Ionescu P, Halsey MJ, Trudell JR. Minimum alveolar concentrations of noble gases, nitrogen, and sulfur hexafluoride in rats: helium and neon as nonimmobilizers (nonanesthetics), 419
- see Trudell JR
- Koch CG, see Capdeville M
- Koch ME, see Ayoub CM
- Kochs E, see Lu H
- Koerschgen M, see Pavlin DJ
- Kohda Y, see Miyabe M
- Kohrs R, Durieux ME. Ketamine: teaching an old drug new tricks, 1186
- Koht A, Moss J. Transient global amnesia: a cause for postanesthetic memory disorder (letter, reply), 980
- Koinig H, Wallner T, Marhofer P, Andel H, Hörauf K, Mayer N. Magnesium sulfate reduces intra- and postoperative analgesic requirements, 206
- Komatsu H, Matsumoto S, Mitsuhata H, Abe K, Toriyabe S. Comparison of patient-controlled epidural analgesia with and without background infusion after gastrectomy, 907
- Kongstad L, see Schou H
- Konrad Ch, Fieber T, Schüpfer G, Gerber H, Müllner G. Comparing the enzyme allergosorbens and coated allergen particle tests for latex allergy: which *in vitro* test should be chosen by an anesthesiologist?, 1389
- Jöhr M. Blockade of the sciatic nerve in the popliteal fossa: a system for

- standardization in children, 1256
- Korman B, Riley RH. Convulsions induced by ropivacaine during interscalene brachial plexus block (letter, reply), 497
- Korttila K, see Hovorka J
- Kosaka Y  
see Hara K  
see Saito Y  
see Sumi M
- Koshizaki M, see Sumi M
- Kost E, see Nguyen AC
- Koyfman L, see Talmor D
- Kozek ME, see Kozek-Langenecker SA
- Kozek-Langenecker SA, Marhofer P, Krenn CG, Glaser C, Kozek ME, Semsroth M. Simulation of an epidural test dose with intravenous isoproterenol in sevoflurane- and halothane-anesthetized children, 549
- Wanzel O, Berger R, Kettner SC, Coraim F. Increased anticoagulation during cardiopulmonary bypass by prostaglandin E<sub>1</sub>, 985
- Krafft P, Nikolic A, Frass M. Esophageal rupture associated with the use of the Combitube (letter), 1457
- Kramer MG, see Howie MB
- Krauth J, see Arndt JO
- Kreienbühl G, see Heidegger T
- Kreimeier U, see Orth VH
- Kreislner NS, Schmit DB. A simple oxygen delivery system for the transportation of intubated patients (letter), 982
- Krenn CG, Schäfer B, Berlakovich GA, Steininger R, Steltzer H, Spiss CK. Detection of graft nonfunction after liver transplantation by assessment of indocyanine green kinetics, 34  
see Kozek-Langenecker SA
- Kristensen J, see Paech M
- Kritzinger M, see Tschernko EM
- Krock JL, Padda GS, Moore JD. Ohmeda 7900 ventilator flow sensor failure (letter), 231
- Ku PKM, Tong MCF, Ho KM, Kwan A, van Hasselt CA. Traumatic esophageal perforation resulting from endotracheal intubation, 730
- Kubo A, see Kikumoto K
- Kucuk N, Kizilkaya M, Tokdemir M. Preoperative epidural ketamine does not have a postoperative opioid sparing effect, 103
- Kumazawa T, see Iwashita H
- Kundra P, Deepalakshmi K, Ravishankar M. Preemptive caudal bupivacaine and morphine for postoperative analgesia in children, 52
- Knimoto F. Intrapulmonary shunt fraction and prostaglandin E<sub>1</sub> (letter, reply), 233
- Kurehara K, see Sha K
- Kuro M, see Miyashita T
- Kuroda M, see Saff GN
- Kwan A, see Ku PKM
- LaGorio J, Thompson VA, Sternberg D, Dorje P. Akathisia and anesthesia: refusal of surgery after the administration of metoclopramide, 224
- Laippala P, see Puura AIE
- LaQuaglia MP, see Borchardt RA
- Larsson A, see Schou H
- Laster MJ  
see Gong D  
see Koblin DD
- Laterre P-F, see Von Montigny S
- Latto IP, see Asai T
- Le Cras AE, Galley HF, Webster NR. Spinal but not general anesthesia increases the ratio of T helper 1 to T helper 2 cell subsets in patients undergoing transurethral resection of the prostate, 1421
- Le Normand L, see Malinovsky J-M
- Lecarpentier Y, see Lejay M
- Lee A, see Done ML
- Lee DH, see Abdi S
- Lee HT, see Joshi S
- Lee JH, see Tsai PS
- Lee S-C, see Wang J-J
- Lee TC, see Tsai PS
- Leftheriotis G, see Preckel MP
- Leighton BL  
see Fogel ST  
see Huffnagle SL
- Lejay M, Hanouz J-L, Lecarpentier Y, Coriat P, Riou B. Modifications of the inotropic responses to  $\alpha$ - and  $\beta$ -adrenoceptor stimulation by propofol in rat myocardium, 277
- Lema G, Canessa R. Phenylephrine, dopamine, mannitol, and renal protection during cardiopulmonary bypass (letter, reply), 1459
- Lema MJ, see de Leon-Casasola OA
- Leoni A, see Casati A
- Lepage J-Y, see Malinovsky J-M
- Leslie K, see Wheelahan JM
- Leung JM, Voskanian A, Bellows WH, Pastor D. Automated electrocardiograph ST segment trending monitors: accuracy in detecting myocardial ischemia, 4
- Levin A, Datta S, Camann WR. Intrathecal ropivacaine for labor analgesia: a comparison with bupivacaine, 624
- Levy JH, see Mochizuki T
- Lew TWK, see Chow MYH
- Lewis TJ, see Dunn SM
- Li J, see Sonner JM
- Liam BL, see Chow MYH
- Liaw W-J, see Wang J-J
- Liguori GA, Kahn RL, Gordon J, Gordon MA, Urban MK. The use of metoprolol and glycopyrrolate to prevent hypotensive/bradycardic events during shoulder arthroscopy in the sitting position under interscalene block, 1320
- Lindahl SGE, see Selldén E
- Lineberger CK, see Pisetsky MA
- Lippmann M. Complications of CO<sub>2</sub> flooding the surgical field in open heart surgery: an old technique revisited (letter), 978
- Lissoni A, see Pelosi P
- Litman RS. Critical Care of the Surgical Newborn (book review), 235
- Berkowitz RJ, Ward DS. Defining levels of sedation in pediatric patients (letter, reply), 1454
- Liu M. Neural Blockade in Clinical Anesthesia and Management of Pain. 3rd ed. (book review), 749
- Liu SS, see Allen HW
- Lobato EB, see Johnson RF
- Lockridge O, see Vanlinthout LE
- Lockwood GG, see Shaw ADS
- Loick HM, see Sielenkämper AW
- London ED, see Harukuni I
- Long GT, see Munro AJ
- Lopez F-M, see Ripart J
- LoPresti S, see Runza M
- Lowry DW, Mirakhur RK, McCarthy GJ, Carroll MT, McCourt KC. Neuromuscular effects of rocuronium during sevoflurane, isoflurane, and intravenous anesthesia, 936
- Lowson SM, see Hanson EW
- Lu H, Werner C, Engelhard K, Scholz M, Kochs E. The effects of sevoflurane on cerebral blood flow autoregulation in rats, 854
- Lubarsky DA, DeLong ER. The impact of choice of muscle relaxant on postoperative recovery time (letter), 499  
see Pisetsky MA
- Lucchinetti E, see Zaugg M
- Ludington E, Dexter F. Statistical analysis of total labor pain using the visual analog scale and application to studies of analgesic effectiveness



- during childbirth, 723
- Lumb PD, see Groudine S
- Lurie S, see Reuben SS
- Lustik SJ, Eichelberger JP, Chhibber AK, Bronsther O. Torsade de pointes during orthotopic liver transplantation, 300
- Lyden ER, see Hill GE
- Lydon AM, Boylan JF. Reversibility of parkinsonism-induced acute upper airway obstruction by benztrapine therapy, 975
- Lynch JP, see Renz CL
- Ma D, Sapsed-Byrne SM, Chakrabarti MK, Ridout D, Whitwam JG. Synergism between sevoflurane and intravenous fentanyl on A $\delta$  and C somatosympathetic reflexes in dogs, 211
- Maekawa N, see Mikawa K
- Mahajan RP, Cavill G, Simpson EJ. Reliability of the transient hyperemic response test in detecting changes in cerebral autoregulation induced by the graded variations in end-tidal carbon dioxide, 843
- Malhotra V. Endocrine Disorders and Anesthesia. Vol. 35, No. 4 of International Anesthesiology Clinics (book review), 983
- Malinge M, see Malinovsky J-M
- Malinovsky J-M, Le Normand L, Lepage J-Y, Malinge M, Cozian A, Pinaud M, Buzelin J-M. The urodynamic effects of intravenous opioids and ketoprofen in humans, 456
- Malmgren JA, see Pavlin DJ
- Mamie JC, see Van Elstraete AC
- Mantha S, see Fleisher LA
- Marcel RJ, see Ozelsel TJP
- Marcus MAE, Gogarten W, Buerkle H, Van Aken H. New epidural drugs: *primum non nocere* (letter, reply), 1212
- Marhofer P  
see Koinig H  
see Kozek-Langenecker SA
- Marks RA, see Saff GN
- Marquardt B, see Arndt JO
- Marsch SC, see Erb T
- Martin GR, see Bevan JC
- Martin PH, see Crighton IM
- Martinez-Ruiz R, see Kaynar AM
- Martinot S, see Vedrinne JM
- Martins MA, see Capelozzi M
- Marudhachalam KS, see Puri GD
- Marx GF, see Yun EM
- Mather LE, see Grace RF
- Matsuda T, see Fujise K
- Matsukawa T  
see Iwashita H  
see Nishiyama T
- Matsuki A, see Sato T
- Matsumoto S  
see Fujise K  
see Komatsu H
- Matsuno A, see Kawana S
- Matsuo H, see Kikumoto K
- Mautz DS, see Momota Y
- Mayer N, see Koinig H
- Mayes LC, see Kain ZN
- Maze M, see Poree LR
- Mazor A, see Tverskoy M
- Mazurek AJ, Rae B, Hann S, Kim JI, Castro B, Coté CJ. Rocuronium versus succinylcholine: are they equally effective during rapid-sequence induction of anesthesia?, 1259
- Mazze RI. No evidence of sevoflurane-induced renal injury in volunteers (letter), 230
- McCallum JB, Boban N, Hogan Q, Schmeling WT, Kampine JP, Bosnjak ZJ. The mechanism of  $\alpha_2$ -adrenergic inhibition of sympathetic ganglionic transmission, 503
- McCarthy GJ, see Lowry DW
- McCarthy RJ, see Asokumar B
- McCleane GJ. The cholecystokinin antagonist proglumide enhances the analgesic efficacy of morphine in humans with chronic benign pain, 1117
- McCourt KC, see Lowry DW
- McDowall RH, see Borchardt RA
- McFarlane C, see Sarraf-Yazdi S
- McGrimley L, see Ayoub CM
- McGuinness M, see Schug SA
- McNitt JD, Bode ET, Nelson RE. Long-term pharmaceutical cost reduction using a data management system, 837
- McSweeney TD, see Howie MB
- Mecklem D  
see Brimacombe J  
see Brimacombe JR
- Mehdaoui H, see Van Elstraete AC
- Meier PN, see Walz R
- Meißner A, Weber TP, Van Aken H, Zbieranek K, Rolf N. Clonidine improves recovery from myocardial stunning in conscious chronically instrumented dogs, 1009
- Menon DK, see Navapurkar VU
- Mentzelopoulos SD, Augustatou CG, Papageorgiou EP. Capnography-guided nasotracheal intubation of a patient with a difficult airway and unwanted respiratory depression, 734
- Mersiades M, see Grace RF
- Metge L, see Ripart J
- Metz S. Awareness with recall during general anesthesia (letter), 1459
- Metzner J, see Niv D
- Meyer J, see Sielenkämper AW
- Meyer PG, see Orliaguet GA
- Michna E, see Janfaza DR
- Mikami O, see Fujise K
- Mikat-Stevens M, see Pappas ALS
- Mikawa K, Akamatsu H, Nishina K, Shiga M, Maekawa N, Obara H, Niwa Y. Propofol inhibits human neutrophil functions, 695
- Miller F, see Parmet JL
- Miller KF, Spiekermann BF. A simple way to keep your pen germ-free (letter), 495
- Miller RD, Rampil L, Cohen N. Fewer residents: financial, educational, and practical implications (editorial), 242
- Miller-Hance WC, see Russell IAM
- Milon D, see Benhamou D
- Mima H, see Fukuda K
- Minamino N, see Kikumoto K
- Mirakhur RK, see Lowry DW
- Mitchell MR, see Griffin PR
- Mitsuhashi H  
see Iseki M  
see Komatsu H
- Miura Y, see Sarraf-Yazdi S
- Miyabe M, Kakiuchi Y, Kihara S, Takahashi S, Kohda Y, Sato S, Toyooka H. The plasma concentration of lidocaine's principal metabolite increases during continuous epidural anesthesia in infants and children, 1056
- Miyamoto A, see Kanaya N
- Miyamoto E, see Nagata A
- Miyashita T, Kuro M. Evaluation of platelet function by Sonoclot analysis compared with other hemostatic variables in cardiac surgery, 1228
- Miyazaki T, see Iseki M
- Mizobuchi S, see Yokoyama M
- Moawad A, see Wittels B
- Mochizuki T, Olson PJ, Szlam F, Ramsay JG, Levy JH. Protamine reversal of heparin affects platelet aggregation and activated clotting time after cardiopulmonary bypass, 781
- Möllmann M, see Holst D
- Momota Y, Artru AA, Powers KM, Mautz DS, Ueda Y. Posttreatment with

- propofol terminates lidocaine-induced epileptiform electroencephalogram activity in rabbits: effects on cerebrospinal fluid dynamics, 900
- Monchi M, see Joly L-M
- Mongan PD, Brown RS, Thwaites BK.  
Tranexamic acid and aprotinin reduce postoperative bleeding and transfusions during primary coronary revascularization, 258
- Monk TG, Brecher ME, Goodnough LT.  
Acute normovolemic hemodilution versus predonation: what is the actual cost? (letter, reply), 234
- Moore JD, see Krock JL
- Mori H, see Doi Y
- Mori K, see Fukuda K
- Moriarty DC, see O'Connor PJ
- Morikawa H, see Fukuda K
- Morita S, see Nakata Y
- Mormol JS, see Fogel ST
- Morris R  
see Brimacombe J  
see Brimacombe JR
- Moser D, see Hofbauer R
- Moss J  
see Doenicke AW  
see Koht A  
see Renz CL
- Müller M, Padberg W, Schindler E, Sticher J, Osmer C, Friemann S, Hempelmann G. Renocortical tissue oxygen pressure measurements in patients undergoing living donor kidney transplantation, 474
- Müllner G, see Konrad Ch
- Munro AJ, Long GT, Sleight JW. Nurse-administered subcutaneous morphine is a satisfactory alternative to intravenous patient-controlled analgesia morphine after cardiac surgery, 11
- Murphy RL Jr. T.E.A.L.: The Electronic Anesthesiology Library on CD-ROM 1993-1997 (book review), 1461
- Murray MJ. Sepsis and Multiorgan Failure (book review), 235
- Murray WB, Trojanowski A. A nonpharmacological three-step test for confirmation of correct epidural catheter placement (letter), 1216
- Myles C, see Smith CE
- Nader-Djalal N, Knight PR III, Thusu K, Davidson BA, Holm BA, Johnson KJ, Dandona P. Reactive oxygen species contribute to oxygen-related lung injury after acid aspiration, 127
- Nagata A, Nakao S, Miyamoto E, Inada T, Tooyama I, Kimura H, Shingu K. Propofol inhibits ketamine-induced c-fos expression in the rat posterior cingulate cortex, 1416  
see Fujise K
- Nairn CS, see Allen HW
- Nakabayashi K, see Kawana S
- Nakao S, see Nagata A
- Nakata Y, Goto T, Saito H, Ichinose F, Uezono S, Morita S. The placement of the cuffed oropharyngeal airway with sevoflurane in adults: a comparison with the laryngeal mask airway, 143
- Nakatsuka H, see Yokoyama M
- Namiki A  
see Kanaya N  
see Kawana S
- Naruse R, see Chen L
- Nasrallah G, see Baurain MJ
- Navapurkar VU, Skepper JN, Jones JG, Menon DK. Propofol preserves the viability of isolated rat hepatocyte suspensions under an oxidant stress, 1152
- Nelskylä K, see Hovorka J
- Nelson RE, see McNitt JD
- Nemirovsky A, see Niv D
- Neuvonen PJ, see Palkama VJ
- Newman LM, see Asokumar B
- Ng M, Hastings RH. Successful direct laryngoscopy assisted by posture in a patient with ankylosing spondylitis, 1436
- Nguyen AC, Kost E, Framstad M. Indigo carmine-induced severe hypotension, 1194
- Nicolson SC, see Cook-Sather SD
- Nielsen R, see Callesen T
- Nikolic A, see Krafft P
- Nikolov MP, see Chaney MA
- Niquille M, Van Gessel E, Gamulin Z. Continuous spinal anesthesia for hip surgery in a patient with Shy-Drager syndrome, 396
- Nishikawa T  
see Goyagi T  
see Tanaka M
- Nishina K, see Mikawa K
- Nishiyama T, Hanaoka K. Inorganic fluoride kinetics and renal and hepatic function after repeated sevoflurane anesthesia, 468
- Matsukawa T, Hanaoka K. Safety of a new lightwand device (Trachlight): temperature and histopathological study, 717
- Niv D, Nemirovsky A, Metzner J, Rudick V, Jurna I, Urca G. Antinociceptive effect induced by the combined administration of spinal morphine and systemic buprenorphine, 583
- Niwa Y, see Mikawa K
- Norman PH, see Daley MD
- Norris A, see Jin F
- Norris MC  
see Ferouz F  
see Huffnagle SL
- Nossaman BD, see Kaye AD
- Nozaki-Taguchi N, Yamamoto T.  
Involvement of nitric oxide in peripheral antinociception mediated by  $\kappa$ - and  $\delta$ -opioid receptors, 388
- Obara H, see Mikawa K
- O'Connor BR, Park KW. Complications of CO<sub>2</sub> flooding the surgical field in open heart surgery: an old technique revisited (letter, reply), 979
- O'Connor M, see Yared J-P
- O'Connor PJ, Russell JD, Moriarty DC.  
Anesthetic implications of laryngeal trauma, 1283
- Oh J, Camann W. Severe, acute meningeal irritative reaction after epidural blood patch, 1139
- Ohata H, see Ueda N
- Ohshika H, see Kanaya N
- Okaño T, see Takiguchi T
- Okubo Y, see Takiguchi T
- Okuno S, see Iseki M
- Olkkola KT, see Palkama VJ
- Olson PJ, see Mochizuki T
- Ong BC, see Chow MYH
- Orliaguet GA, Meyer PG, Blanot S, Jarreau M-M, Charron B, Buisson C, Carli PA. Predictive factors of outcome in severely traumatized children, 537
- Orth VH, Rehm M, Thiel M, Kreimeier U, Haller M, Brechtelsbauer H, Finsterer U. First clinical implications of perioperative red cell volume measurement with a nonradioactive marker (sodium fluorescein), 1234
- Osmer C, see Müller M
- Ostapkovich N, see Joshi S
- Ostrom DN, see Schwartz DA
- Ovassapian A, see Rosenblatt WH
- Owens BD, see Allen HW
- Ozaki M, see Iwashita H
- Ozelsel TJP, Hein HAT, Marcel RJ, Rathjen KW, Ramsay MAE, Jackson RW.  
Delayed neurological deficit after total hip arthroplasty, 1209
- Öztürk E, see Babacan A
- Pace NL, see Swenson JD

- Padberg W, see Müller M
- Padda GS, see Krock JL
- Paech M, Kristensen J, Ilett KF.  
Nonsteroidal antiinflammatory  
drugs during lactation (letter), 977
- Pagel PS, Fu JL, Damask MC, Davis RF,  
Samuelson PN, Howie MB, Warltier  
DC. Desflurane and isoflurane  
produce similar alterations in  
systemic and pulmonary  
hemodynamics and arterial  
oxygenation in patients undergoing  
one-lung ventilation during  
thoracotomy, 800  
see Hettrick DA
- Paiva PSO, see Capelozzi M
- Palkama VJ, Isohanni MH, Neuvonen PJ,  
Olkola KT. The effect of  
intravenous and oral fluconazole on  
the pharmacokinetics and  
pharmacodynamics of intravenous  
alfentanil, 190
- Palmer SK. Obstetric Anesthesia and  
Uncommon Disorders (book  
review), 750
- Palmon SC, Kirsch JR, Depper JA, Toung  
TJK. The effect of the prone position  
on pulmonary mechanics is frame-  
dependent, 1175
- Panning B, see Walz R
- Papageorgiou EP, see Mentzelopoulos SD
- Pappas ALS, Sukhani R, Hotaling AJ,  
Mikat-Stevens M, Javorski JJ,  
Donzelli J, Shenoy K. The effect of  
preoperative dexamethasone on the  
immediate and delayed  
postoperative morbidity in children  
undergoing adenotonsillectomy, 57
- Paris A, Scholz J, von Knobelsdorff G,  
Tonner PH, Schulte am Esch J. The  
effect of remifentanyl on cerebral  
blood flow velocity, 569
- Park H-S, see Chung C-J
- Park KW, see O'Connor BR
- Parker BM, see Seidman PA
- Parker RK, see Dunn SM
- Parment JL, Colonna-Romano P, Horrow  
JC, Miller F, Gonzales J, Rosenberg  
H. The laryngeal mask airway  
reliably provides rescue ventilation  
in cases of unanticipated difficult  
tracheal intubation along with  
difficult mask ventilation, 661
- Horrow JC, Berman AT, Miller F, Pharo  
G, Collins L. The incidence of large  
venous emboli during total knee  
arthroplasty without pneumatic  
tourniquet use, 439
- Pascale V, see Ayoub CM
- Pasch T  
see Jamnicki M  
see Zaugg M
- Pastor D, see Leung JM
- Pavlin DJ, Rapp SE, Polissar NL,  
Malmgren JA, Koerschgen M, Keyes  
H. Factors affecting discharge time  
in adult outpatients, 816
- Pavy TJG, Gambling DR, Douglas MJ.  
Combination of diclofenac and  
intrathecal morphine for cesarean  
delivery (letter), 1453
- Pearl RG, see Rodriguez RM
- Pearlstein R, see Sarraf-Yazdi S
- Pedlow S, see Ellis JE
- Pedoto A, see Pelosi P
- Pekar A, see Roytblat L
- Pelosi P, Croci M, Ravagnan I, Tredici S,  
Pedoto A, Lissoni A, Gattinoni L.  
The effects of body mass on lung  
volumes, respiratory mechanics,  
and gas exchange during general  
anesthesia, 654
- Penberthy AJ, see Chow TKF
- Pendeville E, see Singelyn FJ
- Perdrix J-P, see Carry P-Y
- Perez de Sá V, see Schou H
- Perrault DF Jr, see Garry B
- Peterfreund RA, see Dull RO
- Peterson MD, see Thomson IR
- Pharo G, see Parmet JL
- Piccirillo JF, see Waddle JP
- Piedmonte M, see Yared J-P
- Pile-Spellman J, see Joshi S
- Pinaud M, see Malinovsky J-M
- Pinchak AC, see Smith CE
- Pisetsky MA, Lubarsky DA, Capehart BP,  
Lineberger CK, Reves JG. Valuing  
the work performed by  
anesthesiology residents and the  
financial impact on teaching  
hospitals in the United States of a  
reduced anesthesia residency  
program size, 245
- Pisini JV, see Janfaza DR
- Pivalizza EG, Tjia IM, Juneja HS, Cohen  
AM, Duke JH Jr. Elective  
splenectomy in an anemic Jehovah's  
Witness patient with cirrhosis, 529
- Plevak DJ, see Sampathkumar P
- Polissar NL, see Pavlin DJ
- Popovic D, see Jamnicki M
- Poree LR, Guo TZ, Kingery WS, Maze M.  
The analgesic potency of  
dexmedetomidine is enhanced after  
nerve injury: a possible role for  
peripheral  $\alpha_2$ -adrenoceptors, 941
- Power I, see Grace RF
- Powers KM, see Momota Y
- Prado AA, see Carvalho JCA
- Prat-Pradal D, see Ripart J
- Prayssac P, see Godet G
- Preckel B, Schlack W, Thämer V. Enflurane  
and isoflurane, but not halothane,  
protect against myocardial  
reperfusion injury after cardioplegic  
arrest with HTK solution in the  
isolated rat heart, 1221
- Preckel MP, Ferber-Viart C, Leftheriotis G,  
Dubreuil C, Duclaux R, Saumet JL,  
Banssillon V, Granry JC.  
Autoregulation of human inner ear  
blood flow during middle ear  
surgery with propofol or isoflurane  
anesthesia during controlled  
hypotension, 1002
- Preis CA, Hartmann T, Zimpfer M.  
Laryngeal mask airway facilitates  
awake fiberoptic intubation in a  
patient with severe oropharyngeal  
bleeding, 728
- Pretorius M, Rasmussen GE, Holcomb  
GW. Hemodynamic and  
catecholamine responses to a  
laparoscopic adrenalectomy for  
pheochromocytoma in a pediatric  
patient, 1268
- Principe D, see Wittels B
- Prough DS, see Bedell EA
- Punjabi A, see Smith CE
- Puri GD, Marudhachalam KS, Chari P,  
Suri RK. The effect of magnesium  
sulphate on hemodynamics and its  
efficacy in attenuating the response  
to endotracheal intubation in  
patients with coronary artery  
disease, 808
- Puura AIE, Rorarius MGF, Laippala P,  
Baer GA. Does abstinence from  
smoking or a transdermal nicotine  
system influence atracurium-  
induced neuromuscular block?, 430
- Qadir S, see Ayoub CM
- Quintin L, de Kock M, Ghignone M.  
Safety of  $\alpha_2$ -agonists? (letter), 496
- Rabito SF, see Sato T
- Rachinsky M, see Roytblat L
- Rae B, see Mazurek AJ
- Raeder JC, see Aasboe V
- Ramajoli F, De Amici D. Is there a bilateral  
block of the thoracic sympathetic  
chain after unilateral intrapleural  
analgesia?, 360
- Ramakrishnan U, see Emmons S
- Rampil L, see Miller RD
- Ramsay JG, see Mochizuki T
- Ramsay MAE, see Ozelsel TJP

- Randle DW, see Greilich PE
- Ranta SO-V, Hynynen M. Awareness with recall during general anesthesia (letter, reply), 1459
- Rapp SE, see Pavlin DJ
- Rasmussen GE, see Pretorius M
- Rathjen KW, see Ozelsel TJP
- Rathmell JP, Viscomi CM, Ashburn MA. Nonsteroidal antiinflammatory drugs during lactation (letter, reply), 977
- Ravagnan I, see Pelosi P
- Ravishankar M, see Kundra P
- Rehm M, see Orth VH
- Reimer EJ, see Bevan JC
- Renz CL, Thurn JD, Finn HA, Lynch JP, Moss J. Oral antihistamines reduce the side effects from rapid vancomycin infusion, 681
- Reuben SS, Connelly NR, Lurie S, Klatt M, Gibson CS. Dose-response of ketorolac as an adjunct to patient-controlled analgesia morphine in patients after spinal fusion surgery, 98
- Steinberg RB, Lurie SD. Is there a place for meperidine in intravenous regional anesthesia? (letter), 1215
- Reves JG, see Pisetsky MA
- Reynolds F, see Russell R
- Rich GF, see Frank DU
- Richardson J, Jones J, Atkinson R. The effect of thoracic paravertebral blockade on intercostal somatosensory evoked potentials, 373
- Richardson MG, Collins HV, Wissler RN. Intrathecal hypobaric versus hyperbaric bupivacaine with morphine for cesarean section, 336
- Ridout D, see Ma D
- Riley RH, see Korman B
- Riopelle JM, see Griffin PR
- Riou B, see Lejay M
- Ripart J, Metge L, Prat-Pradal D, Lopez F-M, Eledjam J-J. Medial canthus single-injection episcleral (sub-tenon anesthesia): computed tomography imaging, 42
- Ririe DG, Butterworth JF, Hines M, Hammon JW Jr, Zaloga GP. Effects of cardiopulmonary bypass and deep hypothermic circulatory arrest on the thyroid axis during and after repair of congenital heart defects: preservation by deep hypothermia?, 543
- Robin J, see Vedrinne JM
- Rockoff MA, Hall SC. Allocating subspecialist pediatric anesthesiologists: when to say no (letter, reply), 742
- Rodriguez RM, Pearl RG. Pulmonary hypertension and major surgery, 812
- Rogers S, see Gan TJ
- Rohling RG, see Biro P
- Rohrich RJ, see Blakeley KR
- Roikjaer O, see Callesen T
- Roizen MF, see Fleisher LA
- Roland P, see Harvey SC
- Rolf N, see Meißner A
- Rorarius MGF, see Puura AIE
- Rose JB, Cohen DE. Pediatric Pain Management and Sedation Handbook (book review), 984
- Rosenberg H, see Parmet JL
- Rosenblatt WH, Wagner PJ, Ovassapian A, Kain ZN. Practice patterns in managing the difficult airway by anesthesiologists in the United States, 153
- Ross AK, Ferrero-Conover D. Anaphylactoid reaction due to the administration of ondansetron in a pediatric neurosurgical patient, 779
- Ross EL, see Janfaza DR
- Rouine-Rapp K, see Russell IAM
- Rowlingson JC. Do we finally have a technique for determining how "blocked" patients are under general anesthesia? (editorial), 1
- Roytblat L, Talmor D, Rachinsky M, Greemberg L, Pekar A, Appelbaum A, Gurman GM, Shapira Y, Duvdenani A. Ketamine attenuates the interleukin-6 response after cardiopulmonary bypass, 266
- see Talmor D
- Rozan JP, see Saff GN
- Rudick V, see Niv D
- Ruetsch YA, see Borgeat A
- Runza M, Albani A, Tagliabue M, Haiek M, LoPresti S, Birnbach DJ. Spinal anesthesia using hyperbaric 0.75% versus hyperbaric 1% bupivacaine for cesarean section, 1099
- Russell IAM, Miller-Hance WC, Silverman NH. Intraoperative transesophageal echocardiography for pediatric patients with congenital heart disease, 1058
- Zwass MS, Fineman JR, Balea M, Rouine-Rapp K, Brook M, Hanley FL, Silverman NH, Cahalan MK. The effects of inhaled nitric oxide on postoperative pulmonary hypertension in infants and children undergoing surgical repair of congenital heart disease, 46
- Russell JD, see O'Connor PJ
- Russell R, Reynolds F. Epidural analgesia and chronic backache (letter), 747
- Saatte S, see Khorasani A
- Sadhasivam S, Subramaniam R. Delleman syndrome: anesthetic implications, 553
- Saff GN, Marks RA, Kuroda M, Rozan JP, Hertz R. Analgesic effect of bupivacaine on extraperitoneal laparoscopic hernia repair, 377
- Saha AK, Higgins M, Walker G, Badr A, Berman L. Comparison of awake endotracheal intubation in patients with cervical spine disease: the lighted intubating stylet versus the fiberoptic bronchoscope, 477
- Saito H, see Nakata Y
- Saito K, see Takiguchi T
- Saito Y, Kaneko M, Kirihaara Y, Sakura S, Kosaka Y. Characteristics of tolerance to somatic and visceral antinociception after continuous epidural infusion of morphine in rats, 1340
- see Hara K
- see Sumi M
- see Tamakawa S
- Saitoh Y, see Fujii Y
- Sakaue G, see Shimaoka M
- Sakura S
- see Hara K
- see Saito Y
- see Sumi M
- Salem MR, see Crystal GJ
- Salfinger H, see Hofbauer R
- Sampathkumar P, Scanlon PD, Plevak DJ. Baclofen withdrawal presenting as multiorgan system failure, 562
- Samuelson PN, see Pagel PS
- Sánchez de la Cuesta F, see De La Cruz JP
- Santos AC, see Yun EM
- Sapsed-Byrne SM
- see Ma D
- see Shaw ADS
- Sarraf-Yazdi S, Sheng H, Miura Y, McFarlane C, Dexter F, Pearlstein R, Warner DS. Relative neuroprotective effects of dizocilpine and isoflurane during focal cerebral ischemia in the rat, 72
- Sarti P, see Xu F
- Sato S, see Miyabe M
- Sato T, Hirota K, Matsuki A, Zsigmond EK, Rabito SF. The role of the N-methyl-D-aspartic acid receptor in the relaxant effect of ketamine on

- tracheal smooth muscle, 1383
- Saumet JL, see Preckel MP
- Sayag A, see Carry P-Y
- Scanlon PD, see Sampathkumar P
- Schäfer B, see Krenn CG
- Schindler E, see Müller M
- Schlack W, see Preckel B
- Schmeling WT, see McCallum JB
- Schmit DB, see Kreisler NS
- Schneider G, see Johansen JW
- Schneider M, see Benhamou D
- Scholz J, see Paris A
- Scholz M, see Lu H
- Schou H, Kongstad L, Perez de Sá V, Werner O, Larsson A. Uncompensated blood loss is not tolerated during acute normovolemic hemodilution in anesthetized pigs, 786
- Schreiner MS. Gastric fluid volume: is it really a risk factor for pulmonary aspiration? (editorial), 754  
see Cook-Sather SD
- Schubert A, see Bloomfield EL
- Schug SA, Sidebotham DA, McGuinnety M, Thomas J, Fox L. Acetaminophen as an adjunct to morphine by patient-controlled analgesia in the management of acute postoperative pain, 368
- Schulte am Esch J, see Paris A
- Schüpfer G, see Konrad Ch
- Schwartz DA, Connelly NR, Theroux CA, Gibson CS, Ostrom DN, Dunn SM, Hirsch BZ, Angelides AG. Gastric contents in children presenting for upper endoscopy, 757
- Scuderi PE, see Tang J
- Sebel PS  
see Bowen PJ  
see Iselin-Chaves IA  
see Johansen JW
- Secic M, see Bloomfield EL
- Sedeño G, see De La Cruz JP
- Segal S, Csavoy AN, Datta S. The tocolytic effect of catecholamines in the gravid rat uterus, 864
- Seidman PA, Parker BM. Sharps disposal in the operating room: current clinical practices and costs, 634
- Seifert B, see Jannicki M
- Selldén E, Lindahl SGE. Amino acid-induced thermogenesis to prevent hypothermia during anesthesia is not associated with increased stress response, 637
- Lindahl SGE. Postoperative nitrogen excretion after amino acid-induced thermogenesis under anesthesia, 641
- Semsroth M, see Kozek-Langenecker SA
- Sessler DI, Badgwell JM. Exposure of postoperative nurses to exhaled anesthetic gases, 1083  
see De Witte JL  
see Iwashita H
- Sha K, Simokawa M, Kawaguchi M, Iwasaka T, Kurehara K, Kitaguchi K, Furuya H. Use of transesophageal echocardiography probe imaging to guide internal jugular vein cannulation, 1032
- Shapira Y  
see Roytblat L  
see Talmor D
- Shapshay SM, see Garry B
- Sharrock NE. Asystole under hypotensive epidural anesthesia (letter), 982
- Shaw ADS, Chamberlain SK, Sapsed-Byrne SM, Lockwood GG. Nitrous oxide and carbon dioxide have no effect on the blood-gas solubilities of sevoflurane and isoflurane, 1412
- Shelley KH, Haddadin AS. Is *Helicobacter pylori* infection an occupational hazard for anesthesiologists?, 973
- Sheng H, see Sarraf-Yazdi S
- Shenoy K, see Pappas ALS
- Sherman SK, see Wang SY
- Shiga M, see Mikawa K
- Shimaoka M, Hosotsubo K, Sugimoto M, Sakaue G, Taenaka N, Yoshiya I, Kiyono H. The influence of surgical stress on T cells: enhancement of early phase lymphocyte activation, 1431
- Shimizu T, see Abe K
- Shimoda O, Ikuta Y, Terasaki H. Assessing the level of regional blockade under general anesthesia using the skin vasomotor reflex test, 83
- Shimonaka H, see Ueda N
- Shingu K  
see Asai T  
see Fujise K  
see Nagata A
- Shiozaki H, see Abe K
- Shoda T, see Fukuda K
- Shutway F, see Bloomfield EL
- Shyken JM, see Fogel ST
- Sidebotham DA, see Schug SA
- Sielenkämper AW, Meyer J, Loick HM, Hachenberg T. Thoracoscopic interruption of patent ductus arteriosus compromises cardiopulmonary function in newborn pigs, 1037
- Sigl J, see Iselin-Chaves IA
- Silbert BS, see Wheelahan JM
- Silverman DG, see Ayoub CM
- Silverman NH, see Russell IAM
- Simokawa M, see Sha K
- Simpson EJ, see Mahajan RP
- Singelyn FJ, Deyaert M, Joris D, Pendeville E, Gouverneur JM. Effects of intravenous patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous three-in-one block on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty, 88
- Sinha PK, Dubey PK, Kaushik S. A simple modification in laryngoscope for continuous suction (letter), 500
- Skepper JN, see Navapurkar VU
- Sleigh JW, see Munro AJ
- Slogoff S, see Chaney MA
- Sloninsky A, see Chen L
- Smeltzer JS, see Fogel ST
- Smiler BG, Srock M. Isopropyl alcohol for transport-related nausea (letter), 1214
- Smith CE, Gerdes E, Sweda S, Myles C, Punjabi A, Pinchak AC, Hagen JF. Warming intravenous fluids reduces perioperative hypothermia in women undergoing ambulatory gynecological surgery, 37
- Smith I, see Ashworth J
- Smith MF, see Bevan JC
- Sniadach MS, Alberts MS. A comparison of the antiemetic effect of ondansetron and droperidol on patients undergoing gynecologic laparoscopy (letter, reply), 1213
- Sollevi A, see von Heijne M
- Somasunderam K, see Thiagarajan A
- Song D, van Vlymen J, White PF. Is the bispectral index useful in predicting fast-track eligibility after ambulatory anesthesia with propofol and desflurane?, 1245
- Sonner JM, Li J, Eger EI II. Desflurane and the nonimmobilizer 1,2-dichlorohexafluorocyclobutane suppress learning by a mechanism independent of the level of unconditioned stimulation, 200
- Li J, Kandel L, Eger EI II. Ethanol concentrations approaching minimum alveolar anesthetic concentration are required to suppress learning in a fear-potentiated startle paradigm in rats, 1398
- Søreide E. Trauma Anesthesia and Critical Care of Neurological Injury (book



- review), 236
- Spahn DR  
  see Jamnicki M  
  see Zaugg M
- Spaulding C, see Joly L-M
- Speer KP, see Klein SM
- Spence JD, see Badner NH
- Spiekermann BF. Subspecialty Care. Vol. 5  
  in *The Atlas of Anesthesia Series*  
  (book review), 1218  
  see Miller KF
- Spiss CK, see Krenn CG
- Sprung J, Castellani WJ, Srinivasan V,  
  Udayashankar S. The effects of  
  donepezil and neostigmine in a  
  patient with unusual  
  pseudocholinesterase activity, 1203
- Srinivasan V, see Sprung J
- Srock M, see Smiler BG
- Starr NJ, see Yared J-P
- Steele SM, Klein SM, D'Ercole FJ,  
  Greengrass RA, Gleason D. A new  
  continuous catheter delivery system  
  (letter), 228  
  see Klein SM
- Steinberg RB  
  see Dunn SM  
  see Reuben SS
- Steininger R, see Krenn CG
- Steltzer H, see Krenn CG
- Sternberg D, see LaGorio J
- Stevenson GW, see Tobin MJ
- Stewart PD, see Crighton IM
- Sticher J, see Müller M
- Suárez E, see Zamacona MK
- Subramaniam R, see Sadhasivam S
- Sugawara K, see Tamakawa S
- Sugimoto M, see Shimaoka M
- Sukhani R, see Pappas ALS
- Sumi M, Sakura S, Koshizaki M, Saito Y,  
  Kosaka Y. The advantages of the  
  lateral decubitus position after  
  spinal anesthesia with hyperbaric  
  tetracaine, 879
- Sun HL. Naloxone: acute opioid  
  withdrawal syndrome or side  
  effects? (letter, reply), 1214
- Sundeman H, Haney M, Broomé M,  
  Häggmark S, Johansson G, Winsö  
  O, Biber B. The effects of desflurane  
  on cardiac function as measured by  
  conductance volumetry in swine,  
  522
- Suri RK, see Puri GD
- Sutin K, see Capan LM
- Suto Y, see Inomata S
- Suwa T, see Inomata S
- Sweda S, see Smith CE
- Swenson JD, Hutchinson DT, Bromberg M,  
  Pace NL. Rapid onset of ulnar nerve  
  dysfunction during transient  
  occlusion of the brachial artery, 677
- Szlam F, see Mochizuki T
- Szmuk P, see see Ezri T
- Taenaka N, see Shimaoka M
- Tagliabue M, see Runza M
- Taittonen MT, Kirvelä OA, Aantaa R,  
  Kanto JH. The effect of clonidine or  
  midazolam premedication on  
  perioperative responses during  
  ketamine anesthesia, 161
- Takada K, see Abe K
- Takahashi S  
  see Fujii Y  
  see Miyabe M
- Takashina M, see Abe K
- Takiguchi T, Okano T, Egawa H, Okubo  
  Y, Saito K, Kitajima T. Epidural  
  saline and combined epidural-spinal  
  anesthesia (letter, reply), 232
- Talmor D, Roytblat L, Artru AA, Yuri O,  
  Koyfman L, Katchko L, Shapira Y.  
  Phenylephrine-induced  
  hypertension does not improve  
  outcome after closed head trauma  
  in rats, 574  
  see Roytblat L
- Tamakawa S, Sugawara K, Yanagita Y,  
  Saito Y. Occult air leak of an  
  endotracheal tube (letter), 746
- Tanabe Y, see Iseki M
- Tanaka H, see Fujii Y
- Tanaka M, Nishikawa T. Epidural saline  
  and combined epidural-spinal  
  anesthesia (letter), 232  
  see Goyagi T
- Tanaka Y, see Doi Y
- Tang J, D'Angelo R, White PF, Scuderi PE.  
  The efficacy of RS-25259, a long-  
  acting selective 5-HT<sub>3</sub> receptor  
  antagonist, for preventing  
  postoperative nausea and vomiting  
  after hysterectomy procedures, 462  
  see Chen L
- Tang JJ-S, see Wang J-J
- Tarantino F, see Fanelli G
- Taylor SM, see Ahmed HE
- Terasaki H, see Shimoda O
- Terazawa E, see Ueda N
- Terrell RC, see Gong D
- Thämer V, see Preckel B
- Theroux CA, see Schwartz DA
- Theroux MC. Tracheal intubation for  
  awake versus paralyzed infants  
  with pyloric stenosis (letter), 1455
- Thiagarajan A, Goverdhan PD, Chari P,  
  Somasunderam K. The effect of  
  hyperventilation and hyperoxia on  
  cerebral venous oxygen saturation  
  in patients with traumatic brain  
  injury, 850
- Thiel M, see Orth VH
- Thomas J, see Schug SA
- Thomas P, see Bourke DL
- Thompson VA, see LaGorio J
- Thomson IR, Peterson MD, Hudson RJ. A  
  comparison of clonidine with  
  conventional preanesthetic  
  medication in patients undergoing  
  coronary artery bypass grafting, 292
- Thorin D, see Benhamou D
- Thorup J, see Callesen T
- Thurn JD, see Renz CL
- Thusu K, see Nader-Djalal N
- Thwaites BK, see Mongan PD
- Tjia IM, see Pivalizza EG
- Tobin MJ, Stevenson GW, Horn BJ, Chen  
  EH, Hall SC, Coté CJ. A comparison  
  of three modes of ventilation with  
  the use of an adult circle system in  
  an infant lung model, 766
- Tokdemir M, see Kucuk N
- Toledano AY, see Wittels B
- Tomlin MK, see Harvey SC
- Tommasi-Rosso M, see Gil F
- Tong MCF, see Ku PKM
- Tonner PH, see Paris A
- Tooyama I, see Nagata A
- Toriyabe S, see Komatsu H
- Torri G  
  see Casati A  
  see Fanelli G
- Torsher LC. Fiberoptic intubation and  
  children (letter), 495
- Toung TJK, see Palmon SC
- Toyooka H  
  see Fujii Y  
  see Inomata S  
  see Miyabe M
- Tramèr MR, Garnerin P. A difference is a  
  difference if it makes a difference  
  (letter), 1460
- Tran T-V, Fiset P, Varin F.  
  Pharmacokinetics and  
  pharmacodynamics of cisatracurium  
  after a short infusion in patients  
  under propofol anesthesia, 1158
- Traystman RJ, see Harukuni I
- Tredici S, see Pelosi P
- Trojanowski A, see Murray WB
- Trott S, see Blakeley KR
- Trudell JR, Koblin DD, Eger EI II. A  
  molecular description of how noble  
  gases and nitrogen bind to a model  
  site of anesthetic action, 411  
  see Koblin DD
- Truelsen KS, Brock-Utne JG. "Damping" of

- an arterial line: an unlikely cause (letter), 979
- Tsai PS, Buerkle H, Huang LT, Lee TC, Yang LC, Lee JH. Lidocaine concentrations in plasma and cerebrospinal fluid after systemic bolus administration in humans, 601
- Tschernko EM, Klepetko H, Gruber E, Kritzing M, Klimscha W, Jandrasits O, Haider W. Clonidine added to the anesthetic solution enhances analgesia and improves oxygenation after intercostal nerve block for thoracotomy, 107
- Tsubokawa T, Yamamoto K, Kobayashi T. Propofol clearance and distribution volume increase in patients with hyperthyroidism, 195
- Tsuchida H, see Kanaya N
- Tuman KJ, see Asokumar B
- Tung C-S, see Wong C-S
- Tunkel DE, see Greenberg RS
- Turndorf H, see Zhou HH
- Tverskoy M, Braslavsky A, Mazor A, Ferman R, Kissin I. The peripheral effect of fentanyl on postoperative pain, 1121
- Uda R, see Doi Y
- Udayashankar S, see Sprung J
- Ueda N, Dohi S, Akamatsu S, Hamaya Y, Terazawa E, Shimonaka H, Ohata H. Pulmonary arterial and right ventricular responses to prophylactic albumin administration before aortic unclamping during abdominal aortic aneurysmectomy, 1020
- Ueda W. Norepinephrine for cardiopulmonary resuscitation during epidural block (letter), 1216
- Ueda Y, see Momota Y
- Uezono S, see Nakata Y
- Umedaly H, see Grace RF
- Urban MK, see Liguori GA
- Urca G, see Niv D
- Urzua J. Interpretation of nonpulsatile arterial pressure-flow relations during cardiopulmonary bypass (letter, reply), 743
- Van Aken H  
see Camu F  
see Knichwitz G  
see Marcus MAE  
see Meißner A
- van Ee R, Hemrika DJ, de Blok S. Multimodal approach to postoperative pain control in daycase laparoscopic sterilization (letter), 496
- Van Elstraete AC, Mamie JC, Mehdaoui H. Nasotracheal intubation in patients with immobilized cervical spine: a comparison of tracheal tube cuff inflation and fiberoptic bronchoscopy, 400
- Van Gessel E  
see Fournier R  
see Niquille M
- van Hasselt CA, see Ku PKM
- van Huffelen AC, see de Vries JW
- van Vlymen J, see Song D
- Vandam LD. The Genesis of Surgical Anesthesia (book review), 501
- Vanderelst P, see Von Montigny S
- Vang MC, see Joshi S
- Vanlinthout LE, Bartels CF, Lockridge O, Callens K, Booij LH. Prolonged paralysis after a test dose of mivacurium in a patient with atypical serum cholinesterase, 1199
- Varin F, see Tran T-V
- Vedrinne C, see Vedrinne JM
- Vedrinne JM, Curtil A, Martinot S, Vedrinne C, Robin J, Franck M, Champsaur G. The hemodynamic effects of hypoxemia in anesthetized pigs: a comparison between right heart catheter and echocardiography, 21
- Veen A, see Broekema AA
- Viguera A, Welch C, Bigatello L, Drop LJ. Hemodynamic responses to electroconvulsive therapy in a hypertensive patient with end-stage pulmonary fibrosis, 737
- Viscomi CM, see Rathmell JP
- Visser GH, see de Vries JW
- Viswanathan S, see Griffin PR
- Vogel W, see Bräu ME
- von Heijne M, Hao J-X, Yu W, Sollevi A, Xu X-J, Wiesenfeld-Hallin Z. Reduced anti-allodynic effect of the adenosine A<sub>1</sub>-receptor agonist R-phenylisopropyladenosine on repeated intrathecal administration and lack of cross-tolerance with morphine in a rat model of central pain, 1367
- von Knobelsdorff G, see Paris A
- Von Montigny S, Laterre P-F, Vanderelst P, De Kock M. The effects of intraoperative intravenous clonidine on gastric intramucosal Pco<sub>2</sub>, 686
- Voskarian A, see Leung JM
- Waddle JP, Evers AS, Piccirillo JF. Postanesthesia care unit length of stay: quantifying and assessing dependent factors, 628
- Wagner PJ, see Rosenblatt WH
- Wahlander S, see Capan LM
- Wahl CC. "Man-in-the-barrel" syndrome after endoscopic sinus surgery, 1196
- Walker G, see Saha AK
- Wallner T, see Koinig H
- Walter M, see Kasper S-M
- Walz R, Bund M, Meier PN, Panning B. Esophageal rupture associated with the use of the Combitube (letter), 228
- Wang J-J, Ho S-T, Lee S-C, Tang JJ-S, Liaw W-J. Intraarticular triamcinolone acetate for pain control after arthroscopic knee surgery, 1113
- Wang SY, Sherman SK. Preferential effect of nitroglycerin on large microvessels (letter), 747
- Wanzel O, see Kozek-Langenecker SA
- Ward CF. A warning concerning InfusO.R. pump labels for remifentanyl (letter), 1214
- Ward DS, see Litman RS
- Ware PD, see Allen HW
- Wartier DC  
see Hettrick DA  
see Pagel PS
- Warner DO, see Kai T
- Warner DS  
see Klein SM  
see Sarraf-Yazdi S
- Watanabe H, see Kawana S
- Watcha MF, see White PF
- Watts ADJ, Eliasziw M, Gelb AW. Propofol and hyperventilation for the treatment of increased intracranial pressure in rabbits, 564
- Weber S, see Joly L-M
- Weber TP, see Meißner A
- Webster NR, see Le Cras AE
- Weiler JM, see Jensen NF
- Weiskopf RB, see Eger EI II
- Welch C, see Viguera A
- Wender RH, see Chen L
- Wendt M, see Holst D
- Werner C, see Lu H
- Werner O, see Schou H
- Wheelahan JM, Leslie K, Silbert BS. Epidural fentanyl reduces the shivering threshold during epidural lidocaine anesthesia, 587
- White PF, Watcha MF. Reversal of neuromuscular blockade (letter), 494  
see Ahmed HE  
see Avramov MN  
see Blakeley KR  
see Chen L  
see Joshi GP



- see Song D  
see Tang J  
Whitwam JG, see Ma D  
Wiesenfeld-Hallin Z, see von Heijne M  
Williams A, see Harvey SC  
Wilson JL, Brown DL, Wong GY, Ehman RL, Cahill DR. Infraclavicular brachial plexus block: parasagittal anatomy important to the coracoid technique, 870  
Wilson RS, see Borchardt RA  
Windsor A, see Bowen PJ  
Windsor AM, see Johansen JW  
Winsö O, see Sundeman H  
Wissler RN, see Richardson MG  
Wittels B, Faure EAM, Chavez R, Moawad A, Ismail M, Hibbard J, Principe D, Karl L, Toledano AY. Effective analgesia after bilateral tubal ligation, 619  
Wong C-S, Cherng C-H, Tung C-S. Intrathecal administration of excitatory amino acid receptor antagonists or nitric oxide synthase inhibitor reduced autotomy behavior in rats, 605  
Wong GY, see Wilson JL  
Woo P, see Garry B  
Worwag E, Chodak GW. Overnight hospitalization after radical prostatectomy: the impact of two clinical pathways on patient satisfaction, length of hospitalization, and morbidity, 62  
Wu W-H, see Hunt TE  
Wurm WH, see Garry B  
Xu F, Sarti P, Zhang J, Blanck TJJ. Halothane and isoflurane alter calcium dynamics in rat cerebrocortical synaptosomes, 701  
Xu X-J, see von Heijne M  
Yaksh TL, see Eisenach JC  
Yamamoto K, see Tsubokawa T  
Yamamoto S, see Kawana S  
Yamamoto T, see Nozaki-Taguchi N  
Yanagita Y, see Tamakawa S  
Yang LC, see Tsai PS  
Yared J-P, Starr NJ, Hoffman-Hogg L, Bashour CA, Insler SR, O'Connor M, Piedmonte M, Cosgrove DM III. Dexamethasone decreases the incidence of shivering after cardiac surgery: a randomized, double-blind, placebo-controlled study, 795  
Yee LL, see Greilich PE  
Yokoyama M, Mizobuchi S, Nakatsuka H, Hirakawa M. Comparison of plasma lidocaine concentrations after injection of a fixed small volume in the stellate ganglion, the lumbar epidural space, or a single intercostal nerve, 112  
Yoshiya I  
see Abe K  
see Shimaoka M  
Young WL, see Joshi S  
Yu W, see von Heijne M  
Yun EM, Marx GF, Santos AC. The effects of maternal position during induction of combined spinal-epidural anesthesia for cesarean delivery, 614  
Yuri O, see Talmor D  
Zahn PK, Brennan TJ. Intrathecal metabotropic glutamate receptor antagonists do not decrease mechanical hyperalgesia in a rat model of postoperative pain, 1354  
Zahurak M, see Greenberg RS  
Zaloga GP, see Ririe DG  
Zalunardo MP, see Zaugg M  
Zamacona MK, Suárez E, García E, Aguirre C, Calvo R. The significance of lipoproteins in serum binding variations of propofol, 1147  
Zammit A, see Grace RF  
Zaugg M, Lucchinetti E, Zalunardo MP, Zumstein S, Spahn DR, Pasch T, Zollinger A. Substantial changes in arterial blood gases during thorascopic surgery can be missed by conventional intermittent laboratory blood gas analyses, 647  
Zbieranek K, see Meißner A  
Zbuzek VK, see Hunt TE  
Zetlaoui PJ. Is there a place for meperidine in intravenous regional anesthesia? (letter, reply), 1216  
Bouaziz H. Lateral approach to the sciatic nerve in the popliteal fossa, 79  
Zhang J, see Xu F  
Zhou HH, Turndorf H. Hyper- and hypoventilation affects spinal motor neuron excitability during isoflurane anesthesia, 407  
Zhou X, see Crystal GJ  
Zimpfer M, see Preis CA  
Zollinger A  
see Jamnicki M  
see Zaugg M  
Zsigmond EK, see Sato T  
Zumstein S, see Zaugg M  
Zwass MS, see Russell IAM

# Subject Index

- Abdominal surgery**, see Surgery
- Abdominal wall lift**, see Anesthetic techniques
- Acetaminophen**  
see Analgesics  
see Measurement techniques, paracetamol absorption test
- Acetylcholinesterase**, see Enzymes
- Activated clotting time**, see Monitoring
- Acute opioid withdrawal syndrome**, see Complications
- Adenosine**, see Pharmacology
- Adenylate cyclase**, see Enzymes
- Adrenalectomy**, see Surgery
- $\alpha$ -Adrenoceptors**, see Receptors
- $\beta$ -Adrenoceptors**, see Receptors
- Adrenomedullin**, see Peptides
- Adult respiratory distress syndrome**, see Lung
- Airway**  
compromise, C1 esterase inhibitor deficiency, 87:480  
cough, smoking history, emergence from isoflurane anesthesia, 87:1170  
difficult  
management, practice patterns by anesthesiologists in United States, 87:153  
management, use of Combitube, patient with cervical spine fracture, 87:1441  
patient with, capnography-guided nasotracheal intubation, 87:734  
emergency techniques, use of LMA  
cases of difficult tracheal intubation along with difficult mask ventilation, 87:661  
facilitation of awake fiberoptic intubation, patient with severe oropharyngeal bleeding, 87:728  
epiglottitis, high-located, large, and proper positioning of LMA and COPA, 87:489  
oropharyngeal distance, assessment, use of MRI, 87:1048
- Airway obstruction**, see Complications
- Airway smooth muscle**, see Muscles
- Akathisia**, see Complications
- Albumin**, see Fluid balance
- Albuterol**, see Pharmacology
- Alfentanil**  
see Analgesics  
see Anesthetics, intravenous  
see Pharmacology
- Allergy**  
latex  
enzyme allergosorbents test vs coated allergen particle test, 87:1389  
patient undergoing heart transplantation, 87:304  
ondansetron, anaphylactoid reaction, pediatric neurosurgical patient, 87:779
- Alprostadil**, see Prostaglandin E<sub>1</sub>
- Alzheimer's disease**, see Complications
- Ambulatory anesthesia**, see Anesthesia
- Amino acids**  
see Nutrition, total peripheral  
see Receptors
- Amitriptyline**, see Analgesics
- Amnesia**, see Complications, transient global amnesia
- Amniotic fluid embolism**, see Embolism
- Analgesics**  
acetaminophen, as adjunct to morphine, PCA, management of acute postoperative pain, 87:368  
alfentanil, and propofol, interaction, effect on recall, loss of consciousness, and Bispectral Index, 87:949  
amitriptyline, anti-allodynic effects, rat model of neuropathic pain, 87:1360  
buprenorphine  
intravenous, vs intravenous ketoprofen, urodynamic effects, humans, 87:456  
systemic, and spinal morphine, combined administration, antinociceptive effect induced by, 87:583  
clonidine  
added to anesthetic solution, effects on analgesia and oxygenation after intercostal nerve block for thoracotomy, 87:107  
intrathecal, vs intravenous, reduction of experimental thermal or capsaicin-induced pain and hyperalgesia, 87:591  
codeine, intravenous, vs intravenous morphine and intravenous tramadol, effects on gastric emptying, human volunteers, 87:445  
diclofenac, combined with intrathecal morphine, for cesarean delivery (letter), 87:1453  
fentanyl  
demand-size doses, administered by PCA, postoperative analgesic effects, 87:890  
during labor, effects of intrathecal bupivacaine, 87:1309  
effect on electrophysiologic recovery of CA 1 pyramidal cells from anoxia, rat hippocampal slice model, 87:68  
eltanolone and, interaction, with special reference to logistic regression analysis, 87:967  
epidural, effect on shivering threshold during epidural lidocaine anesthesia, 87:587  
intrathecal clonidine and fentanyl with hyperbaric bupivacaine, effect on analgesia during cesarean section, 87:609  
intravenous, and sevoflurane, synergism between, on A $\delta$  and C somatosympathetic reflexes, dogs, 87:211  
intravenous, vs intravenous ketoprofen, urodynamic effects, humans, 87:456  
requirements, postoperative, reduction, and perioperative magnesium sulfate administration, 87:206  
gabapentin, anti-allodynic effects, rat model of neuropathic pain, 87:1360  
hydromorphone, postoperative analgesic requirement, effect of location of TENS, acupoint vs nonacupoint stimulation, 87:1129  
indomethacin, during lactation (letter), 87:977  
ketoprofen, intravenous, vs intravenous opioids, urodynamic effects, humans, 87:456  
ketorolac, and morphine, intravenous PCA  
after spinal fusion surgery, dose-response, 87:98  
after total knee replacement surgery, 87:93  
lidocaine, anti-allodynic effects, rat model of neuropathic pain, 87:1360  
meperidine  
analgesia after bilateral tubal ligation, 87:619  
intravenous regional anesthesia (letter), 87:1215

- postoperative analgesic consumption,  
failure of preoperative epidural  
ketamine to decrease, 87:103
- morphine  
acetaminophen as adjunct to, PCA,  
management of acute postoperative  
pain, 87:368  
analgesic efficacy, effect of  
proglumide, humans with chronic  
benign pain, 87:1117  
bupivacaine and, caudal, preemptive,  
for postoperative analgesia,  
children, 87:52  
bupivacaine with, intrathecal,  
hyperbaric vs hypobaric, for  
cesarean section, 87:336  
continuous epidural infusion,  
characteristics of tolerance to  
somatic and visceral  
antinociception, rats, 87:1340  
epidural, with bupivacaine, vs  
epidural sufentanil with  
bupivacaine and intramuscular  
morphine at fixed rate,  
postoperative analgesia, patients  
undergoing major abdominal  
surgery, 87:1346  
intramuscular, at fixed rate, vs  
epidural morphine with  
bupivacaine and epidural sufentanil  
with bupivacaine, postoperative  
analgesia, patients undergoing  
major abdominal surgery,  
87:1346  
intrathecal, combined with diclofenac,  
for cesarean delivery (letter),  
87:1453  
intrathecal, combined with intrathecal  
neostigmine, efficacy, post-cesarean  
section analgesia, 87:341  
intrathecal, efficacy, post-cesarean  
section analgesia, 87:341  
intravenous, vs intravenous codeine  
and intravenous tramadol, effects  
on gastric emptying, human  
volunteers, 87:445  
intravenous, vs intravenous  
ketoprofen, urodynamic effects,  
humans, 87:456  
intravenous PCA, nurse-administered  
subcutaneous morphine as  
satisfactory alternative, 87:11  
intravenous PCA, vs continuous  
epidural analgesia and continuous  
three-in-one block, effects on  
postoperative pain and knee  
rehabilitation after unilateral total  
knee arthroplasty, 87:88
- ketamine as adjunct to,  
postthoracotomy anesthesia,  
unintended N-of-1 study, 87:1372  
ketorolac and, intravenous PCA, after  
spinal fusion surgery, dose-  
response, 87:98  
ketorolac and, intravenous PCA, after  
total knee replacement surgery,  
87:93  
requirements, intraoperative, effect of  
preoperative dextromethorphan,  
patients undergoing laparotomy,  
87:1135  
R-phenylisopropyladenosine and, lack  
of cross-tolerance, rat model of  
central pain, 87:1367  
spinal, and systemic buprenorphine,  
combined administration,  
antinociceptive effect induced by,  
87:583  
subcutaneous, nurse-administered, as  
satisfactory alternative to  
intravenous PCA morphine, 87:11  
nalbuphine, intravenous, vs intravenous  
ketoprofen, urodynamic effects,  
humans, 87:456  
neostigmine, intrathecal, and intrathecal  
neostigmine combined with  
intrathecal morphine, post-cesarean  
analgesia, 87:341
- remifentanyl  
effect on CBF velocity, 87:569  
infusion, intraoperative, complicated  
by infiltration of intravenous  
catheter, followed by recrudescence  
of opioid effects (letter), 87:1458  
warning concerning InfusO.R. pump  
labels (letter), 87:1214
- sufentanil  
epidural, with bupivacaine, vs  
epidural morphine with  
bupivacaine and intramuscular  
morphine at fixed rate,  
postoperative analgesia, patients  
undergoing major abdominal  
surgery, 87:1346  
intrathecal, mechanism of analgesic  
action (letter), 87:1211  
intrathecal, respiratory depression  
after 5  $\mu$ g, 87:1377  
intrathecal, vs epidural lidocaine with  
epinephrine and sufentanil, for  
early labor analgesia, 86:331
- tramadol  
intravenous, vs intravenous codeine  
and intravenous morphine, effects  
on gastric emptying, human  
volunteers, 87:445
- reduction of sweating,  
vasoconstriction, and shivering  
thresholds, 87:173
- Anaphylactoid reaction, see**  
Complications
- Anesthesia**  
ambulatory  
Bispectral Index and fast-track  
eligibility, 87:1245  
desflurane in spontaneously breathing  
patients, 87:312  
discharge time and ambulatory  
anesthesia, 87:816  
drinking fluids after ambulatory  
anesthesia, 87:306  
effect of betamethasone on PONV,  
87:319  
effect of video on recall, 87:531  
episcleral medial canthus block CT  
imaging, 87:42  
preventing hypothermia in  
outpatients, 87:37  
TIVA technique for facial laser  
resurfacing, 87:827
- cardiovascular  
 $\alpha$ -adrenergic inhibition of sympathetic  
ganglia, 87:503  
autoregulation, inner ear blood flow,  
and controlled hypotension, 87:1002  
brain activity and defibrillation  
threshold testing, 87:16  
bronchial blocker with double-lumen  
tube, 87:1239  
cardiac effects of NO, 87:994  
carotid endarterectomy, mandibular  
nerve block, 87:1034  
clonidine, sympathetic nervous  
system, and myocardial stunning,  
87:109  
clonidine vs conventional  
premedication, 87:292  
coagulation, HES 130/0.4 vs HES 200/  
0.5, 87:989  
desflurane and cardiac function,  
87:522  
desflurane and one-lung ventilation,  
87:800  
dexamethasone, shivering, and cardiac  
surgery, 87:795  
echocardiographic assessment of  
hypoxemia, 87:21  
elective splenectomy, Jehovah's  
Witness patient, 87:529  
electrocardiographic ST trending  
monitors and ischemia, 87:4  
graft nonfunction after liver  
transplantation and indocyanine  
green kinetics, 87:34  
hemodilution and blood loss, 87:786

**Anesthesia—continued**

hemodilution with bovine hemoglobin, 87:284  
 internal jugular vein cannulation, use of TEE, 87:1032  
 isoflurane and biventricular afterload, 87:511  
 ketamine and IL-6 response after CPB, 87:266  
 latex allergy, heart transplant patient, 87:304  
 magnesium, intubation, and cardiovascular response, 87:808  
 methylprednisolone and early extubation, 87:27  
 platelet function measured by Sonoclot analysis, 87:1228  
 proinflammatory cytokines, ketamine, and cAMP, 87:1015  
 propofol and myocardial adrenoceptor stimulation, 87:277  
 prostaglandin E<sub>1</sub> and CPB, 87:985  
 protamine, heparin, platelet aggregation, and ACT, 87:781  
 protection against myocardial reperfusion injury, 87:1221  
 pulmonary hypertension and major surgery, 87:812  
 RCV measurement with sodium fluorescein, 87:1234  
 response to albumin during aortic resection, 87:1021  
 sedation after aortic surgery, 87:272  
 spinal cord stimulation for anginal pain, 87:1242  
 subcutaneous vs intravenous PCA morphine, 87:11  
 torsade de pointes with liver transplant, 87:300  
 tranexamic acid and CPB, 87:258  
 water-tight bronchial seal, 87:1027  
 critical care and trauma  
   baclofen withdrawal as multiorgan system failure, 87:562  
   effect of prolonged inhaled NO, 87:1285  
   immobilized cervical spine and intubation, 87:400  
   intramucosal PCO<sub>2</sub> measurement, 87:134  
   laryngeal trauma, anesthetic implications, 87:1283  
   milrinone and diaphragmatic fatigue, 87:1078  
   oxidants in hyperoxia-aspiration lung injury, 87:127  
   pain and arterial cannulation, 87:403  
   percutaneous dilational tracheostomy, 87:556

electroconvulsive therapy, hemodynamic responses, hypertensive patient with end-stage pulmonary fibrosis, 87:737  
 neurosurgical  
   adenosine and cerebrovascular resistance, 87:1291  
   adrenomedullin and subarachnoid hemorrhage, 87:859  
   CO<sub>2</sub> and transient hyperemic response, 87:843  
   continuous CSF drainage, flow-regulated pump, 87:1306  
   effect of fentanyl on anoxic damage, 87:68  
   hyper-/hypoventilation and H reflex/F wave, 87:407  
   hyperventilation, hyperoxia, and brain injury, 87:850  
   induced hypertension and head trauma outcome, 87:574  
   NMDA antagonism and focal cerebral ischemia, 87:72  
   neuroprotection from focal ischemia by PPBP, 87:1299  
   propofol, hyperventilation, and intracranial pressure, 87:564  
   remifentanyl and CBF velocity, 87:569  
   scalp infiltration in craniotomy, 87:579  
   sevoflurane and CBF autoregulation, 87:854  
 obstetric  
   atypical presentation of amniotic fluid embolism, 87:124  
   dystocia risks with labor epidural analgesia, 87:119  
   effective analgesia after tubal ligation, 87:619  
   gynecologic laparoscopy and serotonin excretion, 87:1104  
   hyper- vs hypobaric spinal bupivacaine, 87:336  
   hypotension and CSE anesthesia, 87:614  
   intrathecal clonidine and visceral pain, 87:609  
   intrathecal fentanyl plus bupivacaine, 87:1309  
   intrathecal neostigmine for analgesia, 87:341  
   intrathecal vs epidural in early labor, 87:331  
   labor epidural analgesia technique, 87:326  
   1% vs 0.75% spinal bupivacaine, 87:1099  
   ropivacaine vs bupivacaine for labor analgesia, 87:624  
   tocolytic effect of catecholamines, 87:864

**pediatric**

adult circle system use, infant model, 87:766  
 anaphylaxis and ondansetron administration, 87:779  
 bronchial injury during lung isolation, 87:324  
 bupivacaine-morphine for preemptive analgesia, 87:52  
 continuous epidural anesthesia with lidocaine, 87:1056  
 Delleman syndrome, anesthetic implications, 87:553  
 dexamethasone, children, and adenotonsillectomy, 87:57  
 epidural test dose and isoproterenol, 87:549  
 gastric contents in pediatric endoscopy, 87:757  
 granisetron-droperidol combination, 87:761  
 laparoscopic adrenalectomy for pheochromocytoma, 87:1268  
 mivacurium, children, and sevoflurane, 87:772  
 NO in pediatric surgery, 87:46  
 oropharyngeal distance, assessment using MRI, 87:1048  
 preoperative preparation programs, 87:1249  
 rapid-sequence induction of anesthesia, 87:1259  
 respiratory effects of desflurane, 87:1052  
 risk factors of death in traumatized children, 87:537  
 sciatic nerve blockade in popliteal fossa, 87:1256  
 TEE and congenital heart disease, 87:1058  
 thyroid function during heart surgery, 87:543  
 tracheal extubation and skin incision, sevoflurane MAC, 87:1263  
 VATS for PDA interruption, 87:1037  
 volatile anesthesia in immature hearts, 87:1041  
 regional anesthesia and pain management  
   acetaminophen in postoperative pain control, 87:368  
   antinociception of calcium channel blockers, 87:382  
   background infusion in PCEA after abdominal surgery, 87:907  
   bilateral block and unilateral intrapleural analgesia, 87:360  
   blocking potencies of local anesthetics, 87:885

caudal epidural blood patch for treatment of postdural puncture headache, 87:394  
central pain and R-phenylisopropyladenosine, 87:1367  
clonidine and intercostal blockade, 87:107  
clonidine and pressor response to ephedrine, 87:1336  
combined spinal morphine-systemic buprenorphine, 87:583  
continuous spinal anesthesia for hip surgery, 87:396  
cryoanalgesia for herniorrhaphy, 87:896  
CSF leakage after dural puncture, spinal needles, 87:1331  
dextromethorphan and PCA morphine, 87:1135  
effect of bupivacaine on laparoscopic hernia pain, 87:377  
electrical nerve stimulation and opioid requirement, 87:1129  
epidural fentanyl and shivering, 87:587  
epidural morphine, tolerance characteristics, 87:1340  
excitatory amino acid antagonist, NO synthase inhibitor, and autotomy, 87:605  
facial nerve palsy after scalp block, 87:1365  
fentanyl PCA, 87:890  
fibrin glue patch in an *in vitro* model, 87:1125  
indicator of sensory blockade level, 87:83  
infraclavicular brachial plexus block, 87:870  
interscalene brachial plexus block, 87:1316  
intraarticular triamcinolone for pain control, 87:1113  
intramuscular morphine at fixed rate, 87:1346  
ketamine as adjunct to morphine, 87:1372  
ketorolac, morphine, and PCA, 87:98  
lateral decubitus and hyperbaric spinal, 87:879  
lidocaine concentration after nerve blocks, 87:112  
metabotropic glutamate receptor antagonists, 87:1354  
methicillin-resistant *Staphylococcus aureus* sepsis, 87:116  
neuropathic pain, effects of amitriptyline, gabapentin, and lidocaine, 87:1360

$\kappa$ - and  $\delta$ -opioid receptors and NO, 87:388  
pain relief and knee rehabilitation after total knee arthroplasty, 87:88  
percutaneous electrical nerve stimulation and acute herpes zoster, 87:911  
peripheral effect of fentanyl, 87:1121  
peripheral nerve blocks after total knee replacement, 87:93  
plasma and cerebrospinal lidocaine concentration, 87:601  
preoperative epidural ketamine, for postoperative analgesia, 87:103  
prevention of cardiovascular side effects, spinal anesthesia, 87:347  
progumide increases effects of morphine, 87:1117  
propofol and lidocaine-induced seizures, 87:900  
pulmonary effects of spinal anesthesia, 87:874  
respiratory depression after sufentanil, 87:1377  
ropivacaine for sciatic-femoral block, 87:597  
sciatic nerve block, popliteal fossa, 87:79  
serotonin, nicotine, nifedipine, and analgesia, 87:1109  
shoulder arthroscopy, hypotension/bradycardia, 87:1320  
spinal anesthetics, variations in composition, 87:1326  
spinal vs intravenous clonidine, 87:591  
thoracic paravertebral block and evoked potentials, 87:373  
unilateral spinal anesthesia, 87:355

#### Anesthesiologists

choice of *in vitro* latex allergy test, 87:1389  
occupational hazards, *Helicobacter pylori* infection, 87:973  
pediatric  
allocating subspecialists (letter), 87:742  
subspecialization vs performance-based credentialing (letter), 87:498  
practice patterns in managing difficult airway, 87:153

#### Anesthetics, gases

argon  
binding to sites of anesthetic action, 87:411  
MACs of noble gases, 87:419  
krypton  
binding to sites of anesthetic action, 87:411  
MACs of noble gases, 87:419  
nitrogen

binding to sites of anesthetic action, 87:411  
MACs of noble gases, 87:419  
nitrous oxide  
anesthetic requirements, influence of rat strain, 87:963  
emesis and, correction (letter), 87:1215  
exhaled, exposure of postoperative nurses, 87:1083  
intraoperative use, and perioperative increases in plasma homocysteine, 87:711  
lack of effect on blood-gas solubilities of sevoflurane and isoflurane, 87:1142  
xenon  
binding to sites of anesthetic action, 87:411  
MACs of noble gases, 87:419  
**Anesthetics, intravenous**  
alfentanil, pharmacokinetics and pharmacodynamics, effect of intravenous and oral fluconazole, 87:190  
dexmedetomidine, analgesic potency, enhancement after nerve injury, possible role for peripheral  $\alpha_2$ -adrenoceptors, 87:941  
differential delivery, head and torso, goats, 87:1450  
eltanolone, and fentanyl, interaction, with special reference to logistic regression analysis, 87:967  
fentanyl, peripheral effect on postoperative pain, 87:1121  
ketamine  
as adjunct to morphine, postthoracotomy anesthesia, unintended N-of-1 study, 87:1372  
attenuation of IL-6 response after CPB, 87:266  
effect on proinflammatory cytokine-induced reduction of cardiac intracellular cAMP accumulation, 87:1015  
-induced c-fos expression, in posterior cingulate cortex, effect of propofol, rats, 87:1416  
-induced pulmonary vasodilation, mediation, role of L-type calcium channels, 87:956  
new uses, 87:1186  
perioperative responses during anesthesia, effect of clonidine or midazolam premedication, 87:161  
preoperative epidural, lack of postoperative opioid-sparing effect, 87:103



**Anesthetics, intravenous—continued**

- relaxant effect on tracheal smooth muscle, role of NMDA receptor, 87:1383
- midazolam, vs propofol and sufentanil, sedation after aortic surgery, oxygen consumption and hemodynamic stability, 87:272
- propofol
  - alfentanil and, interaction, effect on recall, loss of consciousness, and Bispectral Index, 87:949
  - ambulatory anesthesia, fast-track eligibility and Bispectral Index, 87:1245
  - anesthesia, neuromuscular effects of rocuronium, 87:936
  - anesthesia, pharmacokinetics and pharmacodynamics of cisatracurium after short infusion, 87:1158
  - clearance and distribution volume increase, patients with hyperthyroidism, 87:195
  - desflurane vs, spontaneously breathing ambulatory patients, 87:312
  - effect on viability of isolated rat hepatocyte suspensions under oxidant stress, 87:1152
  - factors affecting discharge time, adult outpatients, 87:816
  - hyperventilation and, for treatment of increased intracranial pressure, rabbits, 87:564
  - inhibition of human neutrophil functions, 87:695
  - inhibition of ketamine-induced c-fos expression, posterior cingulate cortex, rats, 87:1416
  - in vitro* effects on tissular oxidative stress, rats, 87:1141
  - isoflurane-propofol anesthesia, effects on oxygenation and shunt fraction during one-lung ventilation, 87:1164
  - isoflurane vs, inner ear blood flow autoregulation during controlled hypotension, middle ear surgery, 87:1002
  - isoflurane vs, spontaneously breathing ambulatory patients, 87:312
  - midazolam vs and sufentanil, sedation after aortic surgery, oxygen consumption and hemodynamic stability, 87:272
  - modifications of inotropic responses to  $\alpha$ - and  $\beta$ -adrenoceptor stimulation, rat myocardium, 87:277
  - serum binding variations and lipoproteins, 87:1147

- sevoflurane-propofol anesthesia, effects on oxygenation and shunt fraction during one-lung ventilation, 87:1164

**sufentanil**

- clonidine vs conventional premedication, 87:292
- effect on migration of human leukocytes through human endothelial cell monolayers, 87:1181
- midazolam and propofol vs, sedation after aortic surgery, oxygen consumption and hemodynamic stability, 87:272

**thiopental**

- factors affecting discharge time, adult outpatients, 87:816
- frozen, case (letter), 87:748

**Anesthetics, local****bupivacaine**

- after bilateral tubal ligation, 87:619
- after extraperitoneal laparoscopic hernia repair, 87:377
- epidural morphine or sufentanil with, vs intramuscular morphine at fixed rate, postoperative analgesia, major abdominal surgery, 87:1346
- hyperbaric, with intrathecal clonidine and fentanyl, effect on analgesia during cesarean section, 87:609
- hyperbaric 0.75% vs hyperbaric 1%, spinal anesthesia for cesarean section, 87:1099
- intrathecal, effects on pruritus and duration of fentanyl analgesia during labor, 87:1309
- intrathecal morphine with, hyperbaric vs hypobaric, for cesarean section, 87:336
- mepivacaine vs, sciatic and femoral nerve blockade, 87:597
- morphine and, caudal, for preemptive analgesia, children, 87:52
- ropivacaine vs, labor analgesia, 87:624
- ropivacaine vs, sciatic and femoral nerve blockade, 87:597
- scalp infiltration with, influence on hemodynamics and postoperative pain, adult patients undergoing craniotomy, 87:579
- 0.5% hyperbaric, lateral distribution, effects of spinal needle type, 87:355
- EMLA cream, vs local infiltration anesthesia, for radial artery cannulation, 87:403
- lidocaine
  - anesthesia, epidural, shivering threshold during, effect of epidural fentanyl, 87:587

- concentrations, in plasma and CSF after systemic bolus administration, humans, 87:601
  - epidural, with epinephrine and sufentanil, vs intrathecal sufentanil, for early labor analgesia, 87:331
  - induced epileptiform EEG activity, posttreatment with propofol, effects on CSF dynamics, rabbits, 87:900
  - infusion, use in surgical patient, caution (letter), 87:1212
  - peripheral effect of fentanyl, 87:1121
  - plasma concentrations, after injection of fixed small volume in stellate ganglion, lumbar epidural space, or a single intercostal nerve, comparison, 87:112
  - prilocaine cream, vs local infiltration anesthesia, for radial artery cannulation, 87:403
  - principal metabolite, increased plasma concentration during continuous epidural anesthesia, infants and children, 87:1056
  - mepivacaine, vs bupivacaine and ropivacaine, during sciatic and femoral nerve blockade, 87:597
  - prilocaine, lidocaine-prilocaine cream, vs local infiltration anesthesia, for radial artery cannulation, 87:403
  - ropivacaine
    - bupivacaine vs, labor analgesia, 87:624
    - bupivacaine vs, sciatic and femoral nerve blockade, 87:597
    - comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.75% ropivacaine, interscalene brachial plexus block, 87:1316
    - induced convulsions, during interscalene brachial plexus block (letter), 87:497
    - mepivacaine vs, sciatic and femoral nerve blockade, 87:597
  - tetracaine, hyperbaric, spinal anesthesia with, advantages of lateral decubitus position after, 87:879
- Anesthetics, volatile**
- desflurane**
- ambulatory anesthesia, fast-track eligibility and Bispectral Index, 87:1245
  - anesthesia, respiratory effects on spontaneous ventilation, infants and children, 87:1052
  - anesthetic requirements, influence of rat strain, 87:963
  - effects on cardiac function, as measured by conductance volumetry, swine, 87:522

- exhaled, exposure of postoperative nurses, 87:1083
  - factors affecting discharge time, adult outpatients, 87:816
  - isoflurane vs, effects on systemic and pulmonary hemodynamics and arterial oxygenation, 87:800
  - isoflurane vs, low fresh gas flow rates, control of acute hemodynamic responses during surgery, 87:666
  - isoflurane vs, spontaneously breathing ambulatory patients, 87:312
  - propofol vs, spontaneously breathing ambulatory patients, 87:312
  - suppression of learning by mechanism independent of level of unconditioned stimulation, 87:200
  - enflurane, protective effects against myocardial reperfusion injury after cardioplegic arrest, isolated rat hearts, 87:1221
  - halothane
    - alteration of calcium dynamics, cerebrocortical synaptosomes, rats, 87:701
    - failure to protect against myocardial reperfusion injury after cardioplegic arrest, isolated rat hearts, 87:1221
    - isoflurane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
    - isoflurane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
    - sevoflurane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
    - sevoflurane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
    - sevoflurane vs, simulation of epidural test dose with intravenous isoproterenol, children, 87:549
  - isoflurane
    - alteration of calcium dynamics, cerebrocortical synaptosomes, rats, 87:701
    - anesthesia, emergence from, cough during, 87:1170
    - anesthesia, neuromuscular effects of rocuronium, 87:936
    - anesthesia, spinal motor neuron excitability, effects of hyper- and hypoventilation, 87:407
    - blood-gas solubility, lack of effect of  $N_2O$  and  $CO_2$ , 87:1412
    - clonidine vs conventional premedication, 87:292
    - desflurane vs, effects on systemic and pulmonary hemodynamics and arterial oxygenation, patients undergoing one-lung ventilation during thoracotomy, 87:800
    - desflurane vs, low fresh gas flow rates, control of acute hemodynamic responses during surgery, 87:666
    - desflurane vs, spontaneously breathing ambulatory patients, 87:312
    - disparity of effects on left and right ventricular afterload and hydraulic power generation, swine, 87:511
    - dizocilpine vs, relative neuroprotective effects during focal cerebral ischemia, rats, 87:72
    - exhaled, exposure of postoperative nurses, 87:1083
    - factors affecting discharge time, adult outpatients, 87:816
    - halothane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
    - halothane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
    - MAC, reduction by alfentanil, potentiation by esmolol, 87:671
    - propofol anesthesia, effects on oxygenation and shunt fraction during one-lung ventilation, 87:1164
    - propofol vs, inner ear blood flow autoregulation during controlled hypotension, middle ear surgery, 87:1002
    - propofol vs, spontaneously breathing ambulatory patients, 87:312
    - protective effects against myocardial reperfusion injury after cardioplegic arrest, isolated rat hearts, 87:1221
    - sevoflurane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
    - sevoflurane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
  - sevoflurane
    - and torsade de pointes (letters), 87:981
    - anesthesia, neuromuscular effects of rocuronium, 87:936
    - blood-gas solubility, lack of effect of  $N_2O$  and  $CO_2$ , 87:1412
    - effects on cerebral blood flow autoregulation, rats, 87:854
    - end-tidal concentration, for tracheal extubation and skin incision, children, 87:1263
  - for placement of COPA or LMA, adults, 87:143
  - halothane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
  - halothane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
  - halothane vs, simulation of epidural test dose with intravenous isoproterenol, children, 87:549
  - induced renal injury, lack of evidence in volunteers (letter), 87:230
  - intravenous fentanyl and, synergism between, on A $\delta$  and C somatosympathetic reflexes, dogs, 87:211
  - isoflurane vs, drug-specific effects on  $Ca^{2+}$  sensitization in airway smooth muscle, 87:425
  - isoflurane vs, myocardial depression, cultured neonatal rat ventricular myocytes, 87:1041
  - propofol anesthesia, effects on oxygenation and shunt fraction during one-lung ventilation, 87:1164
  - repeated administration, inorganic fluoride kinetics and renal and hepatic function after, 87:468
- Anesthetic systems, see Equipment**
- Anesthetic techniques**
- abdominal wall lift, effects on respiratory mechanics during laparoscopic cholecystectomy, 87:1393
  - caudal, bupivacaine and morphine, preemptive, for postoperative analgesia in children, 87:52
  - cervical plexus block, plus mandibular nerve block, for carotid endarterectomy, 87:1034
  - CSE
    - cesarean delivery, effects of maternal position, 87:614
    - epidural saline and (letter), 87:232
    - intrathecal vs epidural in early labor, 87:331
    - ropivacaine vs bupivacaine for labor analgesia, 87:624
  - computer-assisted continuous infusion
    - alfentanil, esmolol potentiates isoflurane MAC reduction, 87:671
    - clonidine vs conventional premedication, 87:292
    - interaction of etlanolone and fentanyl, 87:967
  - continuous epidural infusion
    - increased plasma concentration of lidocaine's principal metabolite, infants and children, 87:1056



**Anesthetic techniques—continued**

- intravenous PCA with morphine vs, effects on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty, 87:88
- morphine, characteristics of tolerance to somatic and visceral antinociception, rats, 87:1340
- pain control, patient with herpes zoster, 87:116
- continuous peripheral nerve blockade, new continuous catheter delivery system (letter), 87:228
- continuous spinal anesthesia, for hip surgery, patient with Shy-Drager syndrome, 87:396
- continuous three-in-one block, vs intravenous PCA with morphine, effects on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty, 87:88
- cryoanalgesia, effect on postherniorrhaphy pain, 87:896
- epidural
  - catheter placement, correct, confirmation, nonpharmacological three-step test (letter), 87:1216
  - chronic backache and (letter), 87:747
  - /general anesthesia, assessing level of regional blockade, use of skin vasomotor reflex test, 87:83
  - hypotensive, asystole under (letter), 87:982
  - intravenous PCA vs, rates of myocardial ischemia after aortic surgery (letter), 87:745
  - labor analgesia, and incidence of cesarean delivery for dystocia, 87:119
  - lidocaine anesthesia, shivering threshold during, effect of epidural fentanyl, 87:587
  - needle bevel orientation, influence on spread of sensory blockade, laboring parturient, 87:326
  - test dose, simulation with intravenous isoproterenol, sevoflurane- and halothane-anesthetized children, 87:549
- epidural block
  - cardiopulmonary resuscitation during, use of norepinephrine (letter), 87:1216
  - intercostal nerve block and stellate ganglion block vs, plasma lidocaine concentrations, 87:112
- epidural blood patch
  - acute meningeal irritative reaction, 87:1139

- caudal approach, for treatment of postdural puncture headache, 87:394
- femoral nerve block
  - combined sciatic-femoral nerve block, double-blind comparison of ropivacaine, bupivacaine, and mepivacaine, 87:597
  - improvement of analgesia after total knee replacement surgery, 87:93
- general
  - assessing level of regional blockade, use of skin vasomotor reflex test, 87:83
  - awareness with recall during (letter), 87:1459
  - determining level of neural blockade (editorial), 87:1
  - effect of body mass on lung volumes, respiratory mechanics, and gas exchange, 87:654
  - followed by acute pseudotongue enlargement, 87:1443
  - mechanisms, noble approach (editorial), 87:239
  - spinal vs, ratio of T helper 1 to T helper 2 cell subsets, patients undergoing TURP, 87:1421
- glossopharyngeal nerve block, relief of chronic refractory hiccups (letter), 87:980
- hemodilution
  - acute normovolemic, efficacy (letter), 87:497
  - acute normovolemic, poor tolerance to uncompensated blood loss, anesthetized pigs, 87:786
  - acute normovolemic, vs predonation, actual cost (letter), 87:233
  - preoperative, for elective abdominal aortic surgery, patients undergoing, hemodynamics and oxygen transport, effects of increased doses of bovine hemoglobin, 87:284
  - preoperative, isovolemic, RCV measurement with sodium fluorescein, first clinical implications, 87:1234
- infraclavicular brachial plexus block, parasagittal anatomy important to coracoid technique, 87:870
- intercostal nerve block
  - epidural block and stellate ganglion block vs, plasma lidocaine concentrations, 87:112
  - thoracotomy for, effect of adding clonidine to anesthetic solution, 87:107
- interscalene brachial plexus block
  - comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.75% ropivacaine, 87:1316

- convulsions induced by ropivacaine during (letter), 87:497
- shoulder arthroscopy, in sitting position, use of metoprolol and glycopyrrolate to prevent hypotensive/bradycardic events, 87:1320
- intraarticular steroids, triamcinolone acetate, for pain control after arthroscopic knee surgery, 87:1113
- intrapleural analgesia, unilateral, bilateral block of thoracic sympathetic chain, 87:360
- intravenous regional, use of meperidine (letter), 87:1215
- mandibular nerve block, in addition to cervical plexus block, for carotid endarterectomy, 87:1034
- medial canthus single-injection episcleral (sub-tenon anesthesia), CT imaging, 87:42
- neuromuscular blockade, reversal (letter), 87:494
- one-lung ventilation
  - bronchial injury during lung isolation, pediatric patient, 87:324
  - double-lumen tube and tracheal diameter, 87:158
  - during thoracotomy, systemic and pulmonary hemodynamics and arterial oxygenation, effects of desflurane vs isoflurane, 87:800
  - oxygenation and shunt fraction, effects of propofol, isoflurane, and sevoflurane, 87:1164
  - practical use of bronchial blocker in combination with double-lumen endotracheal tube, 87:1239
  - undetected blood gas changes during, 87:647
- PCA
  - acetaminophen as adjunct to morphine, management of acute postoperative pain, 87:368
  - fentanyl, demand-size doses, postoperative analgesic effects, 87:890
  - intravenous morphine, after cardiac surgery, nurse-administered subcutaneous morphine as alternative, 87:11
  - intravenous morphine, after laparotomy, postoperative requirements, lack of effect of preoperative dextromethorphan, 87:1135
  - intravenous morphine, after spinal fusion surgery, dose-response of ketorolac as adjunct, 87:98

- intravenous morphine, vs continuous epidural analgesia and continuous three-in-one block, effects on postoperative pain and knee rehabilitation, unilateral total knee arthroplasty, 87:88
- intravenous, vs epidural analgesia, rates of myocardial ischemia after aortic surgery (letter), 87:745
- postoperative opioid analgesic requirement, effect of location of TENS, acupoint vs nonacupoint stimulation, 87:1129
- PCEA, with or without background infusion, after gastrectomy, 87:907
- percutaneous electrical nerve stimulation, as alternative to antiviral drugs for acute herpes zoster, 87:911
- preemptive analgesia
  - caudal bupivacaine-morphine, children, 87:52
  - preoperative dextromethorphan, effect on intraoperative morphine requirements, patients undergoing laparotomy, 87:1135
  - preoperative epidural ketamine, lack of postoperative opioid-sparing effect, 87:103
- preperitoneal fascial block, effect of bupivacaine on laparoscopic hernia pain, 87:377
- regional
  - lidocaine concentration after nerve blocks, 87:112
  - of scalp, temporary facial nerve palsy after, 87:1375
- sciatic nerve block
  - addition to femoral nerve block, after total knee replacement surgery, 87:93
  - combined sciatic-femoral nerve block, double-blind comparison of ropivacaine, bupivacaine, and mepivacaine, 87:597
  - popliteal fossa, lateral approach, 86:79
  - popliteal fossa, system for standardization, children, 87:1256
- spinal
  - antinociception of calcium channel blockers, 87:382
  - effect on pulmonary responsiveness to methacholine, guinea pigs, 87:874
  - excitatory amino acid receptor antagonists, NOS inhibitor, and autotomy, 87:605
  - general vs, ratio of T helper 1 to T helper 2 cell subsets, patients undergoing TURP, 87:1421
  - hyperbaric 0.75% vs hyperbaric 1% bupivacaine, cesarean section, 87:1099
  - hyperbaric tetracaine with, advantages of lateral decubitus position after, 87:879
  - incidence and time course of cardiovascular side effects after prophylactic administration of intravenous fluids or vasoconstrictors, 87:347
  - intrathecal clonidine and visceral pain, 87:609
  - intrathecal R-phenylisopropyladenosine, treatment of central pain, rats, 87:1367
  - pressor response to ephedrine, effect of oral clonidine premedication, 87:1336
  - respiratory depression after 5  $\mu$ g intrathecal sufentanil, 87:1377
  - spinal cord stimulator, trial, bedside implantation, for intractable anginal pain, 87:1242
  - stellate ganglion block, vs epidural block and intercostal nerve block, plasma lidocaine concentrations, 87:112
  - subcutaneous, vs topical, radial artery cannulation, 87:403
  - thoracic
    - paravertebral blockade, effect on intercostal somatosensory evoked potentials, 87:373
    - water-tight bronchial seal, 87:1027
  - topical, vs subcutaneous, radial artery cannulation, 87:403
  - total intravenous anesthesia (TIVA)
    - magnesium sulfate and postoperative analgesia, 87:206
    - technique, outpatient facial laser resurfacing, 87:827
  - transcutaneous electrical nerve stimulation (TENS), location, effect on postoperative opioid analgesic requirement, acupoint vs nonacupoint stimulation, 87:1129
- Ankylosing spondylitis**, see Complications
- Anoxia**, see Brain
- Antagonists, heparin**, use after CPB, 87:781
- Antagonists, neuromuscular relaxants**, neostigmine, and donepezil, effects in patient with unusual pseudocholinesterase activity, 87:1203
- Antagonists, opioid**, naloxone
  - acute opioid withdrawal syndrome or side effects (letter), 87:1214
  - partial agonistic activity on opioid receptors expressed from complementary DNAs, Chinese hamster ovary cells, 87:450
- Antibiotics**, vancomycin, rapid infusion, side effects, reduction, effectiveness of oral antihistamines, 87:681
- Antifibrinolytics**
  - aprotinin, vs tranexamic acid, effects on postoperative bleeding and transfusions during primary coronary revascularization, 87:258
  - tranexamic acid, vs aprotinin, effects on postoperative bleeding and transfusions during primary coronary revascularization, 87:258
- Antihistamines**, oral, reduction of side effects from rapid vancomycin infusion, 87:681
- Antimycotics**, fluconazole, intravenous, effect on pharmacodynamics and pharmacokinetics of alfentanil, 87:190
- Antinociception**, see Pain
- Antithrombin III**, see Blood
- Aortic surgery**, see Surgery
- Aprotinin**, see Antifibrinolytics
- Argon**, see Anesthetics, gases
- Arteries**
  - brachial, transient occlusion, rapid onset of ulnar nerve dysfunction during, 87:677
  - coronary, blood flow, hemodilution and blood loss, anesthetized pigs, 87:786
  - radial, cannulation, topical lidocaine-prilocaine cream (EMLA) vs local infiltration anesthesia, 87:403
- Arthroplasty**
  - see Surgery, hip
  - see Surgery, knee
- Arthroscopic surgery**, see Surgery, shoulder
- Asians**, see Ethnic background
- Asthma**, see Complications
- Asystole**, see Complications
- Atracurium**, see Neuromuscular relaxants
- Automated record keeper**, see Measurement techniques
- Autotomy score**, see Measurement techniques
- Awareness**, see Complications
- Backache**, see Complications
- Baclofen**, see Pharmacology
- Benztropine**, see Pharmacology
- Betamethasone**, see Pharmacology
- Bispectral Index**
  - see Anesthetics, intravenous, propofol
  - see Monitoring, electroencephalography

**Blood**

- anticoagulation, increased by
  - prostaglandin E<sub>1</sub> during CPB, 87:985
- antithrombin III, treatment, heparin resistance, and ACT values (letter), 87:1215
- bovine hemoglobin, effects of increased doses on hemodynamics and oxygen transport, patients undergoing preoperative hemodilution for elective abdominal aortic surgery, 87:284
- coagulation
  - compromised, HES 130/0.4 vs HES 200/0.4, assessment, use of thromboelastography, 87:989
  - platelet function, evaluation by Sonoclot analysis vs other hemostatic variables, cardiac surgery, 87:1228
  - protamine, heparin, platelet aggregation, and ACT, 87:781
- leukocytes, human, migration through human endothelial cell monolayers, effect of sufentanil, 87:1181
- loss, uncompensated, poor tolerance during acute normovolemic hemodilution, anesthetized pigs, 87:786
- lymphocytes
  - activation, early phase, enhancement, influence of surgical stress on T cells, 87:1431
  - suppression, postoperative, failure of premedication with clonidine to attenuate, 87:1426
- red cells, volume, measurement, perioperative, with sodium fluorescein, first clinical implications, 87:1234
- transfusion
  - incidence, during primary coronary revascularization, effects of tranexamic acid vs aprotinin, 87:258
  - predonation, vs acute normovolemic hemodilution, actual cost (letter), 87:233

**Blood flow**

- see Brain
- see Ear

**Blood pressure**

- arterial pressure-flow relations, interpretation, during CPB (letter), 87:743
- hypertension
  - induced, after traumatic brain injury (editorial), 87:751

- patient with end-stage pulmonary fibrosis, hemodynamic responses to electroconvulsive therapy, 87:737
- phenylephrine-induced, failure to improve outcome after closed head trauma, rats, 87:574
- hypotension
  - caused by rapid vancomycin infusion, reduction, use of oral antihistamines, 87:681
  - controlled, autoregulation of inner ear blood flow during middle ear surgery with propofol or isoflurane anesthesia, 87:1002
  - CSE anesthesia for cesarean delivery, 87:614
  - prevention of cardiovascular side effects during spinal anesthesia, 87:347
  - severe, indigo carmine-induced, 87:1194
- patients undergoing surgery during ketamine anesthesia, effect of clonidine or midazolam premedication, 87:161
- pressor response to ephedrine, during spinal anesthesia, effect of oral clonidine premedication, 87:1336

**Book and multimedia reviews**

- Atlas of Transesophageal Echocardiography, 87:749
- Clinical Neuroanesthesia. 2nd ed., 87:983
- Clinical Pediatric Anesthesia, 85:236
- Critical Care of the Surgical Newborn, 87:235
- Echocardiography in Pediatric Heart Disease. 2nd ed., 87:1218
- Endocrine Disorders and Anesthesia. Vol. 35, No. 4 of International Anesthesiology Clinics, 87:983
- The Genesis of Surgical Anesthesia, 87:501
- Management of Acute Pain. Vol. 10, No. 1 of Problems in Anesthesia, 87:984
- The Management of Pain, 85:237
- Neural Blockade in Clinical Anesthesia and Management of Pain. 3rd ed., 87:749
- Obstetric Anesthesia and Uncommon Disorders, 87:750
- Pain Relief in Labour. 1st ed., 87:749
- Pediatric Pain Management and Sedation Handbook, 87:984
- Perioperative Transfusion Medicine, 87:235
- Sepsis and Multiorgan Failure, 87:235
- Subspecialty Care. Vol. 5 in The Atlas of Anesthesia Series, 87:1218

T.E.A.L.: The Electronic Anesthesiology Library on CD-ROM 1993-1997, 87:1461

Trauma Anesthesia and Critical Care of Neurological Injury, 85:236

**Bovine hemoglobin**, see Blood

**Bowel function**, postoperative, stimulation (letter), 87:500

**Brachial artery**, see Arteries

**Bradycardia**, see Heart

**Brain**

- anoxia, electrophysiological recovery of CA 1 pyramidal cells, effect of fentanyl, rat hippocampal slice model, 87:68
- blood-brain barrier, propofol and lidocaine-induced seizures, 87:900
- blood flow
  - autoregulation, CO<sub>2</sub> and transient hyperemic response, 87:843
  - autoregulation, effects of sevoflurane, rats, 87:850
  - measurement, initial slope index, assessment of feasibility of manipulating human cerebrovascular resistance with adenosine, 87:1291
  - velocity, effect of remifentanyl, 87:569
- cerebrocortical synaptosomes, calcium dynamics, alteration by halothane and isoflurane, rats, 87:701
- differential delivery of intravenous drugs, goats, 87:1450
- electrical activity, changes during defibrillation threshold testing, 87:16
- intracranial pressure, increased, treatment, use of propofol and hyperventilation, rabbits, 87:564
- ischemia
  - focal, neuroprotection by PPBP, dependence on treatment duration, rats, 87:1299
  - focal, relative neuroprotective effects of dizocilpine and isoflurane, rats, 87:72
  - hyperventilation, hyperoxia, and brain injury, 87:850
- microemboli, delayed neurological deficit after total hip arthroplasty, 87:1209
- microvessels, large, preferential effect of nitroglycerin (letter), 87:747
- oxygen uptake, changes during defibrillation threshold testing, 87:16
- posterior cingulate cortex, ketamine-induced c-fos expression, effect of propofol, rats, 87:1416

- somatosensory evoked potentials,
  - intercostal, effect of thoracic paravertebral blockade, 87:373
- venous oxygen saturation, effect of hyperventilation and hyperoxia, patients with traumatic brain injury, 87:850
- Brain injury**, see Complications, trauma
- Bronchial blocker**, see Equipment
- Bronchial injury**, see Complications
- Bronchoscope**, see Equipment
- Bupivacaine**, see Anesthetics, local
- Buprenorphine**, see Analgesics
- C1 esterase inhibitor deficiency**, see Complications
- CABG**, see Surgery, coronary artery bypass grafting
- Calcium**, see Ions
- Capnography**, see Monitoring
- Capsaicin**, -induced pain and hyperalgesia, effects of intrathecal vs intravenous clonidine, normal volunteers, 87:591
- Carbon dioxide**
  - elimination, efficiency during high-frequency jet ventilation for laryngeal microsurgery, 87:180
  - flooding surgical field during open heart surgery, complications (letter), 87:978
  - lack of effect on blood-gas solubilities of sevoflurane and isoflurane, 87:1142
- Cardiac arrest**, see Complications
- Cardioplegic arrest**, see Heart
- Cardiopulmonary bypass**, see Surgery
- Cardiovascular anesthesia**, see Anesthesia
- Carotid endarterectomy**, see Surgery
- Catecholamines**, see Sympathetic nervous system
- Catheters**, see Equipment
- Caudal anesthesia**, see Anesthetic techniques
- Cells**, endothelial, monolayers, migration of leukocytes through, effect of sufenatril, 87:1181
- Central venous line**, see Equipment
- Cerebral autoregulation**, see Monitoring
- Cerebrospinal fluid (CSF)**
  - dynamics, effects of propofol as treatment for lidocaine-induced epileptiform EEG activity, rabbits, 87:900
  - leaks
    - after dural puncture with various spinal needles, *in vitro* investigation, 87:1331
    - after intracranial procedures, controlled continuous spinal drainage, use of flow-regulated pump, 87:1306
  - lidocaine concentrations after systemic bolus administration, humans, 87:601
- Cervical plexus block**, see Anesthetic techniques
- Cesarean section**, see Surgery
- c-fos**, see Proteins
- Cholecystokinin**, see Neuropeptides
- Cholesterol**, serum protein binding of propofol, 87:1147
- Cholinesterase**, see Enzymes
- Cimetidine**, see Pharmacology
- Cirrhosis**, see Liver
- Cisatracurium**, see Neuromuscular relaxants
- Clonidine**
  - see Analgesics
  - see Pharmacology
  - see Premedication
- Coagulation**, see Blood
- Coagulopathy**, see Complications
- Codeine**, see Analgesics
- Combined spinal-epidural**, see Anesthetic techniques
- Combitube**, see Equipment
- Complications**
  - acute opioid withdrawal syndrome, naloxone (letter), 87:1214
  - airway obstruction
    - caused by pharyngeal and laryngeal edema, succinylcholine and (letter), 87:229
    - jet ventilation, new device, description and model lung testing, 87:915
    - parkinsonism-induced, reversal, benzotropine therapy, 87:975
  - akathisia, after metoclopramide administration, followed by refusal of surgery, 87:224
  - Alzheimer's disease, effects of donepezil and neostigmine, patient with unusual pseudocholinesterase activity, 87:1203
  - anaphylactoid reaction, due to ondansetron administration, pediatric neurosurgical patient, 87:779
  - ankylosing spondylitis, patient with, successful direct laryngoscopy assisted by posture, 87:1436
  - asthma, pulmonary effects of spinal anesthesia, 87:874
  - asystole, hypotensive epidural anesthesia (letter), 87:982
  - awareness, with recall, during general anesthesia (letter), 87:1459
  - backache, chronic, and epidural analgesia (letter), 87:747
  - bleeding
    - oropharyngeal, severe, patient with, awake fiberoptic intubation, facilitation with LMA, 87:728
    - postoperative, after primary coronary revascularization, tranexamic acid vs aprotinin, 87:258
    - prostaglandin E<sub>1</sub> and CPB, 87:985
    - protamine, heparin, platelet aggregation, and ACT, 87:781
  - bradycardia, during shoulder arthroscopy in sitting position under interscalene block, prevention, use of metoprolol and glycopyrrolate, 87:1320
  - bronchial injury, during lung isolation, pediatric patient, 87:324
  - C1 esterase inhibitor deficiency, airway compromise, and anesthesia, 87:480
  - cardiac arrest, norepinephrine for cardiopulmonary resuscitation during epidural block (letter), 87:1216
  - cervical spine disease, awake endotracheal intubation, lighted intubating stylet vs fiberoptic bronchoscope, 87:477
  - chronic hypoxia, effect of prolonged inhaled NO, rats, 87:1285
  - coagulopathy, atypical presentation of amniotic fluid embolism, 87:124
  - convulsions, induced by ropirocaine, during interscalene brachial plexus block (letter), 87:497
  - coronary artery disease, patients with, response to endotracheal intubation, attenuating, efficacy of magnesium sulphate, 87:808
  - coughing, after tracheal intubation, reduction, use of topical steroids, 87:714
  - Dellemann syndrome, anesthetic implications, 87:553
  - diabetes mellitus, patient with, methicillin-resistant *Staphylococcus aureus* sepsis, 87:116
  - dural puncture
    - CSF leakage, *in vitro* investigation, various spinal needles, 87:1331
    - postdural puncture headache, treatment, use of caudal epidural blood patch, 87:394
    - postdural puncture leakage, *in vitro* model, effect of fibrin glue patch, 87:1125
    - unintended, labor epidural analgesia technique, 87:326

**Complications—continued**

- dystocia, cesarean delivery for, incidence, epidural labor analgesia and, 87:119
- Eisenmenger's syndrome, pulmonary hypertension and major surgery, 87:812
- esophageal perforation, traumatic, resulting from endotracheal intubation, 87:730
- esophageal rupture, associated with use of Combitube (letters), 87:228, 87:1457
- facial nerve palsy, temporary, after scalp block, 87:1375
- gastric aspiration, upper endoscopy, children, 87:757
- headache, postdural puncture, treatment caudal epidural blood patch, 87:394 fibrin glue patch, 87:1125
- heart defects, congenital, repair, thyroid axis, effects of CPB and deep hypothermic circulatory arrest, children, 87:543
- heart disease, congenital infants and children undergoing surgical repair, effects of inhaled NO on pulmonary hypertension, 87:46
- intraoperative TEE, pediatric patients, 87:1058
- herpes zoster acute, percutaneous electrical nerve stimulation as alternative to antiviral drugs, 87:911
- patient with, continuous epidural infusion for pain control, methicillin-resistant *Staphylococcus aureus* sepsis resulting from infection of paravertebral muscle, 87:116
- hiccups, chronic refractory, relief with glossopharyngeal nerve block (letter), 87:980
- hoarseness, after tracheal intubation, reduction, use of topical steroids, 87:714
- hyperkalemia, caused by succinylcholine, effects of verapamil and diltiazem pretreatment, dogs, 87:691
- hyperthyroidism, patients with, propofol clearance and distribution volume increase, 87:195
- hypotension, during shoulder arthroscopy in sitting position under interscalene block, prevention, use of metoprolol and glycopyrrolate, 87:1320
- ischemia, upper extremities, and postoperative ulnar neuropathy, 87:677
- Jehovah's Witness, anemic patient with cirrhosis, elective splenectomy, 87:529
- laryngeal stenosis, jet ventilation, new device, 87:915
- "man-in-the-barrel" syndrome, after endoscopic sinus surgery, 87:1196
- meningeal irritative reaction, after epidural blood patch, 87:1139
- mucosal pressure, with LMA, anesthetized adult patients, 87:1379
- multiorgan system failure, baclofen withdrawal presenting as, 87:562
- muscle fasciculations, succinylcholine-induced, prevention, comparison of rocuronium, *d*-tubocurarine, and "mini-dose" succinylcholine, 87:719
- nausea, transport-related, isopropyl alcohol for (letter), 87:1214
- neurological deficit, delayed, after total hip arthroplasty, 87:1209
- neuropathy, ulnar, postoperative, caused by transient occlusion of brachial artery, 87:677
- obesity, effect on lung volumes, respiratory mechanics, and gas exchange during general anesthesia, 87:654
- parkinsonism, -induced acute upper airway obstruction, reversal, use of benztropine therapy, 87:975
- pheochromocytoma, laparoscopic adrenalectomy, hemodynamic and catecholamine responses, pediatric patient, 87:1268
- prolonged neuromuscular block after test dose of mivacurium, patient with atypical serum cholinesterase, 87:1199
- effects of donepezil and neostigmine, patient with unusual pseudocholinesterase activity, 87:1203
- pruritus, associated with fentanyl alone vs intrathecal fentanyl plus bupivacaine, 87:1309
- pseudotongue enlargement, acute, after general anesthesia, 87:1443
- pulmonary aspiration, gastric fluid volume as risk factor (editorial), 87:754
- pulmonary embolism emboli during total knee arthroplasty, 87:439
- massive intraoperative, coincident with succinylcholine administration, 87:491
- pulmonary fibrosis, hypertensive patient with, hemodynamic responses to electroconvulsive therapy, 87:737
- pyloric stenosis, infants with, awake vs paralyzed tracheal intubations (letters), 87:1455
- "red man syndrome," caused by rapid vancomycin infusion, reduction, use of oral antihistamines, 87:681
- respiratory depression after 5  $\mu$ g intrathecal sufentanil, 87:1377
- unwanted, capnography-guided nasotracheal intubation, 87:734
- Shy-Drager syndrome, patient with, continuous spinal anesthesia for hip surgery, 87:396
- sialoadenitis, acute sublingual, after general anesthesia, 87:1443
- sialoadenopathy, acute transient, and LMA, 87:1438
- smoking, and atracurium-induced neuromuscular block, 87:430
- sore throat, after tracheal intubation, reduction, use of topical steroids, 87:714
- subarachnoid hemorrhage, patients with, increased plasma concentration of adrenomedullin, 87:859
- torsade de pointes during, but not necessarily caused by, sevoflurane anesthesia (letter), 87:981
- during orthotopic liver transplantation, 87:300
- sevoflurane and (letter), 87:981
- transdermal nicotine system, and atracurium-induced neuromuscular block, 87:430
- transient global amnesia, cause for postanesthetic memory disorder (letter), 87:980
- trauma brain injury, effect of hyperventilation and hyperoxia on cerebral venous oxygen saturation, 87:850
- brain injury, induced hypertension after (editorial), 87:751
- cervical spine fracture, Combitube elective anesthetic airway management, 87:1441
- facial, patient with severe oropharyngeal bleeding, awake fiberoptic intubation, facilitation with LMA, 87:728



- head, closed, failure of phenylephrine-induced hypertension to improve outcome, rats, 87:574
- laryngeal, anesthetic implications, 87:1283
- risk factors associated with death, children, 87:537
- Computed tomography**, see Measurement techniques
- Computer applications**, data management, and drug cost reduction, 87:837
- Computer-assisted continuous infusion**, see Anesthetic techniques
- Conductance volumetry**, see Measurement techniques
- Consultation**, cardiology, preoperative, intended purpose and perceived utility, survey, 87:830
- Continuous epidural infusion**, see Anesthetic techniques
- Continuous peripheral nerve blockade**, see Anesthetic techniques
- Continuous spinal anesthesia**, see Anesthetic techniques
- Continuous three-in-one block**, see Anesthetic techniques
- Convulsants**
  - 1,2-dichlorohexafluorocyclobutane, convulsant activity, influence of rat strain, 87:963
  - difluoromethyl-1-chlorotetrafluoroethyl ether, convulsant activity, influence of rat strain, 87:963
  - flurothyl, convulsant activity, influence of rat strain, 87:963
- Coronary artery bypass grafting (CABG)**, see Surgery
- Coronary artery disease**, see Complications
- Cost**, see Economics
- Coughing**
  - see Complications
  - see Recovery
- CPB**, see Surgery, cardiopulmonary bypass
- Craniotomy**, see Surgery
- Critical care and trauma**, see Anesthesia
- Cryoanalgesia**, see Anesthetic techniques
- CSE**, see Anesthetic techniques, combined spinal-epidural
- Cuffed oropharyngeal airway (COPA)**, see Equipment
- Cystometry**, see Measurement techniques
- Cytokines**
  - proinflammatory, -induced reduction of cardiac intracellular cAMP accumulation, effect of ketamine, 87:1015
  - ketamine and IL-6 response after CPB, 87:266
- Data management**, see Computer applications
- Death**, see Outcome
- Delleman syndrome**, see Complications
- Desflurane**, see Anesthetics, volatile
- Dexamethasone**, see Pharmacology
- Dexmedetomidine**
  - see Anesthetics, intravenous
  - see Pharmacology
- Dextromethorphan**, see Pharmacology
- Diabetes mellitus**, see Complications
- Diaphragmatic fatigue**, see Muscle
- 1,2-Dichlorohexafluorocyclobutane**
  - see Convulsants
  - see Nonimmobilizers
- Diclofenac**, see Analgesics
- Difluoromethyl-1-chlorotetrafluoroethyl ether**, see Convulsants
- Dihydroergotamine**, see Pharmacology
- Diltiazem**, see Pharmacology
- Diphenhydramine**, see Pharmacology
- Dizocilpine**, see Pharmacology
- Dobutamine**, see Pharmacology
- Domperidone**, see Pharmacology
- Donepezil**, see Pharmacology
- Dopamine**, see Pharmacology
- Dorsal horn**, see Spinal cord
- Droperidol**, see Pharmacology
- Dural puncture**, see Complications
- Dyes**
  - indigo carmine, -induced severe hypotension, 87:1194
  - indocyanine green, detection of graft nonfunction after liver transplantation, 86:34
- Dystocia**, see Complications
- Ear**, inner ear, blood flow, autoregulation, during middle ear surgery with propofol or isoflurane anesthesia during controlled hypotension, 87:1002
- Echocardiography**, see Monitoring
- Economics**
  - and health systems research
  - cardiology consult survey, 87:830
  - data management and drug cost reduction, 87:837
  - incremental cost of substitution, 87:245
  - level of training and dental injury, 87:255
  - measuring patient satisfaction with anesthetic care, review of current methodology, 87:1089
  - medical technology assessment, overview, 87:1271
  - PACU length-of-stay, 87:628
- PACU nursing exposure to exhaled anesthetic gases**, 87:1083
- postprostatectomy hospitalization**, 87:62
- sharps disposal in the OR**, 87:634
- cost**
  - actual, acute normovolemic hemodilution vs predonation (letter), 87:233
  - analgesic effectiveness during childbirth, and statistical analysis of total labor pain using visual analog scale, 87:723
  - implications of fewer residents, 86:242
  - incremental cost of substitution, 87:245
  - reduction, long-term pharmaceutical, using data management system, 87:837
- Edema**
  - see Larynx
  - see Pharynx
- Education**
  - patients
    - preoperative preparation programs in children, comparative examination, 87:1249
    - video, use to convey preanesthetic information, patients undergoing ambulatory surgery, 87:531
  - residents
    - fewer, financial, educational, and practical implications, 87:242
    - incremental cost of substitution, 87:245
- Eisenmenger's syndrome**, see Complications
- Electrocardiography**, see Monitoring
- Electroconvulsive therapy**, see Anesthesia
- Electroencephalography (EEG)**
  - see Measurement techniques
  - see Monitoring
- Eltanolone**, see Anesthetics, intravenous
- Embolism**
  - amniotic fluid, atypical presentation, 87:124
  - pulmonary
    - emboli during total knee arthroplasty, 87:439
    - massive intraoperative, coincident with succinylcholine administration, 87:491
    - thromboembolism, atrial fibrillation and, 87:217
- EMLA cream**, see Anesthetics, local
- Endoscopy**, see Surgery
- Enflurane**, see Anesthetics, volatile
- Enzymes**
  - acetylcholinesterase, inhibitor, donepezil, and neostigmine, effects, patient with unusual pseudocholinesterase activity, 87:1203

**Enzymes—continued**

- adenylate cyclase, partial agonistic activity of naloxone, 87:450
- cholinesterase
  - plasma, mivacurium, children, and sevoflurane, 87:772
  - serum, atypical, patient with, prolonged paralysis after test dose of mivacurium, 87:1199
- GTPase, partial agonistic activity of naloxone, 87:450
- pseudocholinesterase, activity, unusual, patient with, effects of donepezil and neostigmine, 87:1203

**Ephedrine**, see Pharmacology**Epidemiology**, see Statistics**Epidural anesthesia**, see Anesthetic techniques**Epidural blood patch**, see Anesthetic techniques**Epiglottitis**, see Airway**Epinephrine**, see Sympathetic nervous system, catecholamines**Equipment**

- anesthetic systems, semi-closed circle, infant lung model, 87:766
- arterial line, "damping" of, unlikely cause (letter), 87:979
- bronchial blocker, in combination with double-lumen endotracheal tube, practical use, 87:1239
- bronchoscope, fiberoptic, vs lighted intubating stylet, awake endotracheal intubation, cervical spine disease, 87:477
- catheters
  - continuous, new delivery system (letter), 87:228
  - endotracheal tube exchange, double-to single-lumen, facilitation, use of plastic rod/sleeve combination, patients with difficult glottic visualization (letter), 87:744
  - epidural, confirmation of correct placement, nonpharmacological three-step test (letter), 87:1216
  - intravenous, infiltration, recrudescence of opioid effects after intraoperative remifentanyl infusion (letter), 87:1458
  - right heart catheter, vs echocardiography, hemodynamic effects of hypoxemia, anesthetized pigs, 87:21
  - central venous line, failed, interluminal connection (letter), 87:744
- Combitube
  - associated esophageal rupture (letters), 87:228, 87:1457

- elective anesthetic airway management, patient with cervical spine fracture, 87:1441
- cuffed oropharyngeal airway (COPA)
  - difficult positioning, due to high, large epiglottis, 87:489
  - LMA vs, in anesthetized adult patients, 87:147
  - LMA vs, placement, use of sevoflurane, adults, 87:143
- endobronchial tube, double-lumen
  - left, bronchial cuff inflation, comparison of techniques, water-tight bronchial seal, 87:1027
  - predicting size based on tracheal diameter, 87:158
- endotracheal tube
  - double-lumen, in combination with bronchial blocker, 87:1239
  - exchange catheter, double- to single-lumen, facilitation, use of plastic rod/sleeve combination, patients with difficult glottic visualization (letter), 87:744
  - LMA as alternative, ambulatory anesthesia (letter), 87:977
  - occult air leak (letter), 87:746
  - oxygen delivery system, for transportation of intubated patients (letter), 87:982
  - tracheal cuff, method to protect (letter), 87:1217
- epidural infusion pump, flow-regulated, for controlled continuous spinal drainage, 87:1306
- filter needle and filter straw, variations in composition of spinal anesthetic solutions, effects of drug addition order and preparation methods, 87:1326
- fluid warmer, Hotline, reduction of perioperative hypothermia, ambulatory gynecological surgery, 87:37
- frames, surgical, effect of prone position on pulmonary mechanics, 87:1175
- infusion system pumps, InfusO.R. pump labels, for remifentanyl, warning (letter), 87:1214
- intubating laryngeal mask, tracheal intubation through, use of lighted stylet (letter), 87:979
- laryngeal mask airway (LMA)
  - as alternative to tracheal tube, use during ambulatory anesthesia (letter), 87:977
  - associated acute transient sialoadenopathy, 87:1438

- comparison of pharyngeal mucosal pressure and airway sealing pressure, anesthetized adult patients, 87:1379
- COPA vs, in anesthetized adult patients, 87:147
- COPA vs, placement, use of sevoflurane, adults, 87:143
- disposable vs reusable, paralyzed adult patients, 87:921
- facilitation of awake fiberoptic intubation, patient with severe oropharyngeal bleeding, 87:728
- proper positioning, disturbed, due to high, large epiglottis, 87:489
- rescue ventilation, cases of unanticipated difficult tracheal intubation and difficult mask ventilation, 87:661
- use of lighted stylet, tracheal intubation (letter), 87:498
- laryngoscope
  - Bullard Intubating Laryngoscope, tips for users, 87:1206
  - simple modification for continuous suction (letter), 87:500
- lighted stylet
  - fiberoptic bronchoscope vs, awake endotracheal intubation, cervical spine disease, 87:477
  - LMA and, tracheal intubation (letter), 87:498
  - new lightwand device (Trachlight), safety, temperature and histopathological study, 87:717
  - use for tracheal intubation through intubating LMA (letter), 87:979
- needles, spinal
  - CSF leakage after dural puncture, *in vitro* investigation, 87:1331
  - lateral distribution of 0.5% hyperbaric bupivacaine, 87:355
- Ohmeda 7900 Ventilator Flow Sensor, failure (letter), 87:231
- pens, protection from contamination (letter), 87:495
- syringes, prefilled, availability, obstetric units and ORs, 87:1457
- total laryngeal bypass device, description and model lung testing, 87:915
- Univent tubes, damaged, 87:732
- Esmolol**, see Pharmacology
- Esophageal rupture**, see Complications
- Ethanol**, concentrations required to suppress learning in fear-potentiated startle paradigm, rats, 87:1398



**Ethnic background**, Asians, double-lumen tube and tracheal diameter, 87:158

**Extubation**, tracheal

early, CABG patients, pulmonary effects of methylprednisolone, 87:27

end-tidal sevoflurane concentration, children, 87:1263

**Eye**, episcleral medial canthus block CT imaging, 87:42

**Facial laser resurfacing**, see Surgery

**Facial nerve palsy**, see Complications

**Fast-track eligibility**, see Recovery

**Fear conditioning**

effect of ethanol on learning, 87:1398

nonimmobilizers, anesthetics, and learning, 87:200

**Femoral nerve block**, see Anesthetic techniques

**Fentanyl**

see Analgesics

see Anesthetics, intravenous

**Fibrin glue**, see Pharmacology

**Filter needle and filter straw**, see Equipment

**Fluconazole**, see Antimycotics

**Fluid balance**

albumin, administration, prophylactic, before aortic unclamping during abdominal aortic aneurysmectomy, pulmonary and right ventricular responses, 87:1020

prevention of cardiovascular side effects during spinal anesthesia, 87:347

**Fluid warmer**, see Equipment

**Fluids**, drinking, before discharge from ambulatory surgery, 87:306

**Fluoride**, see Ions

**Flurothyl**, see Convulsants

**Formalin test**, see Measurement techniques

**Frames, surgical**, see Equipment

**Free radicals**

antioxidant reserve, oxidants in hyperoxia-aspiration lung injury, 87:127

oxidant injury, isolated rat hepatocyte suspensions, protective effect of propofol, 87:1152

propofol and tissular oxidative stress, rats, 87:1141

**Gabapentin**, see Analgesics

**Gas chromatography**, see Measurement techniques

**Gas exchange**, during general anesthesia, effect of body mass, 87:654

**Gases, noble**

binding to model site of anesthetic action, molecular description, 87:411

mechanisms of anesthetic action

(editorial), 87:239

**Gases, nonanesthetic**, nitric oxide (NO)

influence on vascular, metabolic, and contractile responses to dobutamine, *in situ* canine hearts, 87:994

inhaled, effects on postoperative

pulmonary hypertension, infants and children undergoing surgical repair of congenital heart disease, 87:46

involvement in peripheral

antinociception mediated by  $\kappa$ - and  $\delta$ -opioid receptors, 87:388

prolonged inhalation, effect on

pulmonary vasoconstriction, rats, 87:1285

pulmonary hypertension and major surgery, 87:812

synthase (NOS), inhibitor, intrathecal administration, effect on autotomy behavior, rats, 87:605

**Gastrectomy**, see Surgery

**Gastric aspiration**, see Complications

**Gastric tonometry**, see Measurement techniques

**Gastrointestinal monitoring**, see Monitoring

**Gastrointestinal tract**

gastric emptying, comparison of effects of intravenous tramadol, codeine, and morphine, human volunteers, 87:445

gastric fluid volume, as risk factor for pulmonary aspiration (editorial), 87:754

mucosal perfusion, effects of intraoperative intravenous clonidine, 87:686

**General anesthesia**, see Anesthetic techniques

**Glossopharyngeal nerve block**, see Anesthetic techniques

**Glycopyrrolate**, see Pharmacology

**Granisetron**, see Pharmacology

**GTPase**, see Enzymes

**Gynecological surgery**, see Surgery

**Hallux valgus repair**, see Surgery

**Halothane**, see Anesthetics, volatile

**Health care delivery**

implications of fewer residents, 86:242  
incremental cost of substitution, 87:245

**Heart**

arrhythmia, torsade de pointes, during orthotopic liver transplantation, 87:300

atrial fibrillation, and thromboembolism, 87:217

bradycardia, prevention of

cardiovascular side effects during spinal anesthesia, 87:347

cardioplegic arrest, followed by

myocardial reperfusion injury, protective effects of enflurane and isoflurane, isolated rat hearts, 87:1221

contractility

propofol and myocardial adrenoceptor stimulation, 87:277

responses to dobutamine, influence of NO, *in situ* canine hearts, 87:994

coronary artery blood flow

hemodilution and blood loss, anesthetized pigs, 87:786

responses to dobutamine, influence of NO, *in situ* canine hearts, 87:994

defibrillation testing, changes in cerebral oxygen uptake and electrical activity, 87:16

function, effects of desflurane as

measured by conductance volumetry, swine, 87:522

intracellular cAMP accumulation,

proinflammatory cytokine-induced reduction, inhibition by ketamine, 87:1015

ischemia

after aortic surgery, epidural analgesia vs intravenous PCA (letter), 87:745

detecting, accuracy of automated electrocardiograph ST segment trending monitors, 87:4

myocardial depression, comparative

effects of sevoflurane, isoflurane, and halothane, cultured neonatal rat ventricular myocytes, 87:1041

myocardial oxygen consumption,

responses to dobutamine, influence of NO, *in situ* canine hearts, 87:994

myocardial stunning, recovery from,

effects of clonidine, conscious chronically instrumented dogs, 87:1009

PDA, thoracoscopic interruption, effects

on cardiopulmonary function, newborn pigs, 87:1037

rate

infusion of propofol, sufentanil, or midazolam for sedation after aortic surgery, 87:272

patients undergoing surgery during

ketamine anesthesia, effect of clonidine or midazolam

premedication, 87:161

**Heart—continued**

- reperfusion injury, after cardioplegic arrest, protective effects of enflurane and isoflurane, isolated rat hearts, 87:1221
- tachycardia, ventricular, torsade de pointes, during but not necessarily caused by sevoflurane anesthesia (letter), 87:981
- ventricular function
  - afterload, left and right, disparity of isoflurane effects, swine, 87:511
  - right, response to prophylactic albumin administration before aortic unclamping, abdominal aortic aneurysmectomy, 87:1020

**Heart transplantation**, see Surgery

**Helium**, see Nonimmobilizers

**Hemodilution**, see Anesthetic techniques

**Hemoglobin-based oxygen carrier**, see Oxygen

**Heparin**, see Pharmacology

**Hernia repair**, see Surgery

**Herpes zoster**, see Complications

**Hiccups**, see Complications

**Hip surgery**, see Surgery

**Histamine**, plasma levels, after cisatracurium and vecuronium administration, 87:434

**Homocysteine**, see Metabolism

**Hospitals**, length-of-stay, after radical prostatectomy, impact of two clinical pathways, 87:62

**Hydrogen**, see Nonimmobilizers

**Hydroxyethyl starch**, see Blood, plasma substitutes

**5-Hydroxytryptamine**, see Vasoconstrictors

**Hyperkalemia**, see Complications

**Hyperoxia**, see Oxygen

**Hypertension**

- see Blood pressure
- see Lung

**Hyperthyroidism**, see Complications

**Hypotension**

- see Blood pressure
- see Complications

**Hypothermia**, see Temperature

**Hypothermic circulatory arrest**, deep, effects on thyroid axis before and during repair of congenital heart defects, children, 87:543

**Hypoxemia**, see Oxygen

**Hypoxia**, see Complications, chronic hypoxia

**Hysterectomy**, see Surgery, gynecological

**Indigo carmine**, see Dyes

**Indocyanine green**, see Dyes

**Infection**

- Helicobacter pylori*, as occupational hazard for anesthesiologists, 87:973
- postoperative, influence of surgical stress on T cells, enhancement of early phase lymphocyte activation, 87:1431
- prevention, use of "pen condom" (letter), 87:495
- Staphylococcus aureus*, methicillin-resistant sepsis, resulting from infection in paravertebral muscle after continuous epidural infusion for pain control, patient with herpes zoster, 87:116

**Inflammation**, IL-6, response after CPB, attenuation by ketamine, 87:266

**Infraclavicular brachial plexus block**, see Anesthetic techniques

**Interactions, drug**

- alfentanil
  - fluconazole and, 87:190
  - propofol and, 87:949
- atracurium, and nicotine, 87:430
- donepezil, and neostigmine, 87:1203
- eltanolone, and fentanyl, 87:967
- fentanyl
  - eltanolone and, 87:967
  - lidocaine and, 87:1121
  - sevoflurane and, 87:211
- fluconazole, and alfentanil, 87:190
- isoflurane, and rocuronium, 87:936
- lidocaine, and fentanyl, 87:1121
- neostigmine, and donepezil, 87:1203
- nicotine, and atracurium, 87:430
- propofol
  - alfentanil and, 87:949
  - rocuronium and, 87:936
- rocuronium
  - isoflurane and, 87:936
  - propofol and, 87:936
  - sevoflurane and, 87:936
- sevoflurane
  - fentanyl and, 87:211
  - rocuronium and, 87:936

**Intercostal nerve block**, see Anesthetic techniques

**Interleukin-6**, see Inflammation

**Interscalene brachial plexus block**, see Anesthetic techniques

**Intraarticular steroids**, see Anesthetic techniques

**Intraleural analgesia**, see Anesthetic techniques

**Intravenous regional anesthesia**, see Anesthetic techniques

**Intubation**

- difficult, unanticipated, along with difficult mask ventilation, rescue ventilation provided by LMA, 87:661
  - endotracheal
    - awake fiberoptic, facilitation, use of LMA, patient with severe oropharyngeal bleeding, 87:728
    - awake, lighted intubating stylet vs fiberoptic bronchoscope, patients with cervical spine disease, 87:477
    - conditions, after cisatracurium and vecuronium, 87:434
    - rapid-sequence, effectiveness of rocuronium vs succinylcholine, 87:1259
    - response to, attenuating, efficacy of magnesium sulphate, patients with coronary artery disease, 87:808
    - traumatic esophageal perforation resulting from, 87:730
  - fiberoptic
    - awake, facilitation, use of LMA, patient with severe oropharyngeal bleeding, 87:728
    - children (letter), 87:495
  - nasotracheal
    - capnography-guided, patient with difficult airway and unwanted respiratory depression, 87:734
    - in patients with immobilized cervical spine, tracheal tube cuff inflation vs fiberoptic bronchoscopy, 87:400
    - sore throat, hoarseness, and coughing after, reduction, use of topical steroids, 87:714
  - tracheal
    - assisted by posture, patient with ankylosing spondylitis, 87:1436
    - awake vs paralyzed, infants with pyloric stenosis (letters), 87:1455, 87:1455
    - cough during emergence from isoflurane anesthesia, 87:1170
    - technique, Bullard Intubating Laryngoscope, 87:1206
    - technique, Combitube, patient with cervical spine fracture, 87:1441
    - use of lighted stylet and LMA (letter), 87:498
    - use of lighted stylet through intubating LMA (letter), 87:979
- Ions**
- calcium
    - Ca<sup>2+</sup> sensitization, in airway smooth muscle, drug-specific effects of volatile anesthetics, 87:425

- channels, L-type, agonist, Bay K 8644, volatile anesthesia in immature hearts, 87:1041
- channels, L-type, blockers, antinociceptive effects on visceral and somatic stimuli, rats, 87:382
- channels, L-type, mediation of ketamine-induced pulmonary vasodilation, 87:956
- dynamics, in cerebrocortical synaptosomes, alteration by halothane and isoflurane, rats, 87:701
- fluoride, inorganic, kinetics after repeated sevoflurane anesthesia, 87:468
- potassium
- channels, in peripheral nerve, tonic block, half-maximal blocking concentrations of local anesthetics, 87:855
  - release, after succinylcholine administration, effects of verapamil and diltiazem pretreatment, dogs, 87:691
  - sodium, channels, in peripheral nerve, tonic block, half-maximal blocking concentrations of local anesthetics, 87:855
- Ischemia**
- see Brain
  - see Complications
  - see Heart
- Isoflurane**, see Anesthetics, volatile
- Isopropyl alcohol**, for transport-related nausea (letter), 87:1214
- Isoproterenol**, see Pharmacology
- Jehovah's Witness**, see Complications
- see Liver
- Jet ventilation**, see Ventilation
- Jugular vein**, see Veins
- Ketamine**, see Anesthetics, intravenous
- Ketoprofen**, see Analgesics
- Ketorolac**, see Analgesics
- Kidney**
- function, after repeated sevoflurane administration, 87:468
  - injury, sevoflurane-induced, lack of evidence in volunteers (letter), 87:230
  - renal protection during CPB, use of phenylephrine, dopamine, and mannitol (letter), 87:1458
- Kidney transplantation**, see Surgery
- Knee**
- popliteal fossa
    - blockade of sciatic nerve, system for standardization, children, 87:1256
    - lateral approach to sciatic nerve, 87:79
    - rehabilitation, after unilateral total knee arthroplasty, comparison of three analgesic techniques, 87:88
- Knee surgery**, see Surgery
- Krypton**, see Anesthetics, gases
- Lactation**, nonsteroidal antiinflammatory drugs during (letter), 87:977
- Laryngeal mask airway (LMA)**, see Equipment
- Laryngeal microsurgery**, see Surgery
- Laryngeal stenosis**, see Complications
- Laryngoscope**, see Equipment
- Larynx**
- edema, airway obstruction caused by, succinylcholine and (letter), 87:229
  - trauma, anesthetic implications, 87:1283
- Laser-Doppler flowmetry**, see Measurement techniques
- Lateral position**, see Position
- Latex**, see Allergy
- Learning and memory**, suppression by anesthetics and nonimmobilizers by mechanism independent of level of unconditioned stimuli, 87:200
- resulting from action at lipid (nonpolar) site, 87:1398
- Length-of-stay**
- see Hospitals
  - see Economics
  - see Postanesthesia care unit
- Leukocytes**, see Blood
- Lidocaine**
- see Analgesics
  - see Anesthetics, local
- Lighted stylet**, see Equipment
- Lipoproteins**, see Proteins
- Liver**
- cirrhosis, anemic Jehovah's Witness patient, elective splenectomy, 87:529
  - function, after repeated sevoflurane administration, 87:468
  - isolated rat hepatocyte suspensions, exposure to oxidant injury, protective effect of propofol, 87:1152
- Liver transplantation**, see Surgery
- LMA**, see Equipment, laryngeal mask airway
- Lorazepam**, see Premedication
- Lung**
- adult respiratory distress syndrome, role of reactive oxygen species, 87:127
  - breathing pattern, respiratory effects of desflurane anesthesia, infants and children, 87:1052
  - effects of methylprednisolone, patients undergoing CABG and early extubation, 87:27
  - endothelium, effect of prolonged inhaled NO, rats, 87:1285
  - hemodynamics, alterations produced by desflurane and isoflurane, patients undergoing one-lung ventilation during thoracotomy, 87:800
- hypertension**
- effect of prolonged inhaled NO, rats, 87:1285
  - major surgery, 87:812
  - postoperative, effects of inhaled NO, infants and children undergoing repair of congenital heart disease, 87:46
  - response to prophylactic albumin administration before aortic unclamping, abdominal aortic aneurysmectomy, 87:1020
- hypoxic pulmonary vasoconstriction**, during hypoxemia, right heart catheter vs echocardiography, anesthetized pigs, 87:21
- intrapulmonary shunt fraction**, and prostaglandin E<sub>1</sub> (letter), 87:233
- pulmonary mechanics**, effect of prone positioning, frame-dependence, 87:1175
- vasoconstriction, effect of prolonged inhaled NO, rats, 87:1285
  - vasodilation, ketamine-induced, mediation, role of L-type calcium channels, 87:946
  - volumes, during general anesthesia, and body mass, 87:654
- Lymphocytes**, see Blood
- MAC**, see Potency, anesthetic
- Magnesium sulfate**, see Pharmacology
- Magnetic resonance imaging (MRI)**, see Measurement techniques
- Mandibular nerve block**, see Anesthetic techniques
- "Man-in-the-barrel" syndrome**, see Complications
- Mannitol**, see Pharmacology
- Mastectomy**, see Surgery
- Measurement techniques**
- automated record keeper, data management and drug cost reduction, 87:837
  - autotomy score, intrathecal administration of excitatory amino acid receptor antagonists or NOS inhibitor, 87:605
  - CT, imaging, medial canthus single-injection episcleral (sub-tenon anesthesia), 87:42
  - conductance volumetry, effects of desflurane on cardiac function, swine, 87:522

**Measurement techniques—continued**

- cystometry, effects of intravenous
  - opioids and ketoprofen, 87:456
- EEG, lidocaine-induced epileptiform
  - EEG activity, posttreatment with propofol, effects on CSF dynamics, rabbits, 87:900
- formalin test,  $\kappa$ - and  $\delta$ -opioid receptors and NO, 87:388
- gas chromatography, blood-gas solubilities of sevoflurane and isoflurane, 87:1412
- gastric tonometry, intramucosal  $P_{CO_2}$  determination, 87:134
- laser-Doppler flowmetry
  - inner ear blood flow autoregulation, effect of controlled hypotension, propofol or isoflurane anesthesia, 87:1002
  - sevoflurane and cerebral blood flow autoregulation, 87:854
- local tissue  $PO_2$ , patients undergoing living donor kidney transplantation, 87:474
- MRI, assessment of oropharyngeal distance, 87:1048
- oximetry, buccal, informed use (letter), 87:494
- paracetamol absorption test, intravenous
  - opioids and gastric emptying, 87:445
- RCV, perioperative, with sodium fluorescein, first clinical implications, 87:1234
- skin vasomotor reflex
  - assessing level of regional blockade under general anesthesia, 87:83
  - determining level of neural blockade, patients under general anesthesia (editorial), 87:1
- Sonoclot analysis, evaluation of platelet function, compared with other hemodynamic variables, cardiac surgery 87:1228
- tetanic fade, RF100 Hz and residual curarization, 87:185
- thromboelastography, coagulation assessment, HES 130/0.4 vs HES 200/0.5, 87:989
- train-of-four ratios, RF100 Hz and residual curarization, 87:185
- transcranial Doppler ultrasonography
  - $CO_2$  and transient hyperemic response, 87:843
  - effect of remifentanyl on CBF velocity, 87:569
- urethral profile, effects of intravenous
  - opioids and ketoprofen, 87:456

- visual analog scale, statistical analysis of total labor pain, 87:723

**Mechanisms of analgesic action,**  
intrathecal sufentanil (letter),  
87:1211

**Mechanisms of anesthetic action**

- binding to sites of anesthetic action, 87:411
- blocking potencies of local anesthetics, 87:885
- effect of ethanol on learning, 87:1398
- general anesthesia (editorial), 87:239
- MACs of noble gases, 87:419
- rat strain, anesthesia, and convulsions, 87:963

**Medial canthus single-injection episcleral anesthesia,** see Anesthetic techniques

**Memory,** see Learning and memory

**Memory function,** effect of interaction of propofol and alfentanil, 87:949

**Meperidine,** see Analgesics

**Meta-analysis,** see Statistics

**Metabolism**

- cytochrome P450 enzymes, interaction of fluconazole with alfentanil, 87:190
- homocysteine, plasma, postoperative increases, and use of intraoperative  $N_2O$ , 87:711
- lidocaine, principal metabolite, increased plasma concentration, continuous epidural anesthesia, infants and children, 87:1056
- serotonin, metabolite, 5-hydroxyindoleacetic acid, excretion, women undergoing gynecological laparoscopy, 87:1104

**Metabotropic glutamate receptors,** see Receptors

**Methacholine,** see Pharmacology

**N-Methyl-D-aspartate,** see Receptors

**Methylprednisolone,** see Pharmacology

**Metoclopramide,** see Pharmacology

**Metoprolol,** see Pharmacology

**Meyer-Overton hypothesis,** see Potency, anesthetic

**Midazolam**

- see Anesthetics, intravenous
- see Premedication

**Milrinone,** see Pharmacology

**Minimum alveolar concentration (MAC),**  
see Potency, anesthetic

**Mivacurium,** see Neuromuscular relaxants

**Monitoring**

- activating clotting time (ACT)
  - heparin resistance, antithrombin II treatment, and ACT values (letter), 87:1215

protamine reversal of heparin after CPB, 87:781

blood gas changes, during thoracoscopic surgery, 87:647

capnography, -guided nasotracheal intubation, 87:734

cerebral autoregulation, changes induced by graded variations in end-tidal  $CO_2$ , reliability of transient hyperemic response test, 87:843

echocardiography, vs right heart catheter, assessing hemodynamic effects of hypoxemia, anesthetized pigs, 87:21

electrocardiography, automated ST segment trending monitors, accuracy in detecting myocardial ischemia, 87:4

effect of interaction of propofol and alfentanil, 87:949

predictor of fast-track eligibility after ambulatory anesthesia with propofol and desflurane, 87:1245

gastrointestinal, measurement of intramucosal  $PCO_2$ , 87:134

peripheral nerve stimulation, RF100 Hz and residual curarization, 87:185

transesophageal echocardiography (TEE) emboli during total knee arthroplasty, 87:439

intraoperative, for pediatric patients with congenital heart disease, 87:1058

probe imaging, to guide internal jugular vein cannulation, 87:1032

**Morphine**

see Analgesics

see Premedication

**Multiorgan system failure,** see Complications

**Muscle**

airway smooth

- $Ca^{2+}$  sensitization, drug-specific effects of volatile anesthetics, 87:425
- relaxant effect of ketamine, role of NMDA receptor, 87:1383

cultured neonatal rat ventricular myocytes, comparative myocardial depression of sevoflurane, isoflurane, and halothane, 87:1041

diaphragmatic fatigue, effects of milrinone, dogs, 87:1077

**Muscle fasciculations,** see Complications

**Nalbuphine,** see Analgesics

**Naloxone,** see Antagonists, opioid

**Nausea**

see Complications

see Vomiting

**Neon**, see Nonimmobilizers

**Neostigmine**

- see Analgesics
- see Antagonists, neuromuscular relaxants

**Nerves**

- thoracic sympathetic chain, bilateral block after unilateral intrapleural analgesia, 87:360
- ulnar, dysfunction, rapid onset, during transient occlusion of brachial artery, 87:677

**Neuralgia**, see Pain, postherpetic neuralgia

**Neurological deficit**, see Complications

**Neuromuscular blockade**, see Anesthetic techniques

**Neuromuscular relaxants**

- atracurium
  - induced neuromuscular block, influence of abstinence from smoking or transdermal nicotine system, 87:430
  - RF100 Hz and residual curarization, 87:185
- choice, impact on postoperative recovery time (letter), 87:499
- cisatracurium
  - administration, onset time, endotracheal intubating conditions, and plasma histamine after, 87:434
  - pharmacokinetics and pharmacodynamics after short infusion, patients under propofol anesthesia, 87:1158
- mivacurium
  - requirements, decreased, and delayed neuromuscular recovery during sevoflurane anesthesia, 87:772
  - test dose, prolonged paralysis after, patient with atypical serum cholinesterase, 87:1199
- pancuronium, effects of donepezil and neostigmine, patient with unusual pseudocholinesterase activity, 87:1203
- rocuronium
  - neuromuscular effects during sevoflurane, isoflurane, and intravenous anesthesia, 87:936
  - succinylcholine vs, effectiveness during rapid-sequence induction of anesthesia, 87:1259
  - d*-tubocurarine vs and "mini-dose" succinylcholine, for preventing succinylcholine-induced muscle fasciculations, 87:719

succinylcholine

- airway obstruction caused by pharyngeal and laryngeal edema (letter), 87:229
- effects of donepezil and neostigmine, patient with unusual pseudocholinesterase activity, 87:1203
- induced muscle fasciculations, prevention, comparison of rocuronium, *d*-tubocurarine, and "mini-dose" succinylcholine, 87:719
- massive intraoperative pulmonary embolism coincident with, 87:491
- potassium release after, effects of verapamil and diltiazem pretreatment, dogs, 87:691
- rocuronium vs, effectiveness during rapid-sequence induction of anesthesia, 87:1259
- d*-tubocurarine, vs rocuronium and "mini-dose" succinylcholine, for preventing succinylcholine-induced muscle fasciculations, 87:719
- vecuronium
  - onset time, endotracheal intubating conditions, and plasma histamine after, 87:434
  - RF100 Hz and residual curarization, 87:185

**Neurons**, spinal motor, excitability, hyper- and hypoventilation during isoflurane anesthesia, 87:407

**Neuropathic pain**, see Pain

**Neuropathy**, see Complications

**Neuropeptides**, cholecystokinin,

- antagonists, proglumide, effect on analgesic efficacy of morphine, humans with chronic benign pain, 87:1117

**Neurosurgical anesthesia**, see Anesthesia

**Neurotransmitters**, serotonin, biosynthesis inhibition, effects on nicotine- and nifedipine-induced analgesia, rats, 87:1109

**Neutrophils**, functions, human, inhibition by propofol, 87:695

**Nicardipine**, see Pharmacology

**Nicotine**, see Pharmacology

**Nifedipine**, see Pharmacology

**Nitric oxide (NO)**, see Gases, nonanesthetic

**Nitrogen**, see Anesthetics, gases

**Nitrogen excretion**, urinary, postoperative, after amino-acid-induced thermogenesis under anesthesia, 87:641

**Nitroglycerin**, see Pharmacology

**Nitrous oxide**, see Anesthetics, gases

**NMDA**, see Receptors,

*N*-methyl-D-aspartate

**Noble gases**, see Gases, noble

**Nonimmobilizers**

- 1,2-dichlorohexafluorocyclobutane, suppression of learning by mechanism independent of level of unconditioned stimulation, 87:200

helium

- binding to sites of anesthetic action, 87:411

MACs of noble gases, 87:419

hydrogen

- binding to sites of anesthetic action, 87:411

MACs of noble gases, 87:419

neon

- binding to sites of anesthetic action, 87:411

MACs of noble gases, 87:419

**Nonradioactive markers**, sodium

- fluorescein, perioperative RCV measurement, first clinical implications, 87:1234

**Norepinephrine**, see Sympathetic nervous system, catecholamines

**NOS**, see Gases, nonanesthetic, nitric oxide, synthase

**Nutrition**, total peripheral, amino acids, -induced thermogenesis

- to prevent hypothermia during anesthesia, and stress response, 87:637

under anesthesia, postoperative nitrogen excretion, 87:641

**Obesity**, see Complications

**Obstetric anesthesia**, see Anesthesia

**Occupational exposure**, postoperative nurses, to exhaled anesthetic gases, 87:18

**Ohmeda 7900 Ventilator Flow Sensor**, see Equipment

**Ondansetron**, see Pharmacology

**One-lung ventilation**, see Anesthetic techniques

**Open heart surgery**, see Surgery

**Operating room (OR)**

- Helicobacter pylori* infection, anesthesia occupational hazard, 87:973
- prefilled syringes, availability (letter), 87:1457
- sharps disposal, current clinical practices and costs, 87:634

**Opioid antagonists**, see Antagonists

**Oropharyngeal bleeding**, see Complications, bleeding

**Oropharyngeal distance**, see Airway



**Outcome**

- death, predictive factors, severely traumatized children, 87:537
- duration of hospitalization, after radical prostatectomy, impact of two clinical pathways, 87:62
- patient satisfaction
  - after radical prostatectomy, impact of two clinical pathways, 87:62
  - with anesthesia care, measuring, review of current methodology, 87:1089

**Oximetry**, see Measurement techniques**Oxygen**

- consumption
  - infusion of propofol, sufentanil, or midazolam for sedation after aortic surgery, 87:272
  - myocardial, responses to dobutamine, influence of NO, *in situ* canine hearts, 87:994
  - patients undergoing surgery during ketamine anesthesia, effect of clonidine or midazolam premedication, 87:161
- delivery system, for transportation of intubated patients (letter), 87:982
- hemoglobin-based oxygen carrier, bovine, increased doses, effects on hemodynamics and oxygen transport, patients undergoing preoperative hemodilution for elective abdominal aortic surgery, 87:284
- hyperoxia
  - associated lung injury, after acid aspiration, role of reactive oxygen species, 87:127
  - effect on cerebral venous oxygen saturation, patients with traumatic brain injury, 87:850
- hypoxemia
  - effect on shivering threshold, rabbits anesthetized with 0.2 MAC isoflurane, 87:1408
  - hemodynamic effects, right heart catheter vs echocardiography, anesthetized pigs, 87:21
- mixed venous oxygen saturation, hemodilution and blood loss, anesthetized pigs, 87:786
- transport, effects of increased doses of bovine hemoglobin, patients undergoing preoperative hemodilution for abdominal aortic surgery, 87:284
- uptake, cerebral, changes during defibrillation threshold testing, 87:16

**Oxygenation**

- after intercostal nerve block for thoracotomy, effect of adding clonidine to anesthetic solution, 87:107
- arterial, during one-lung ventilation, effects of propofol, isoflurane, and sevoflurane, 87:1164

**PACU**, see Postanesthesia care unit**Pain**

- anginal, intractable, bedside implantation of trial spinal cord stimulator, 87:1242
- antinociception
  - effect of combined spinal morphine-systemic buprenorphine, 87:583
  - effects of intrathecal L-type calcium channel blockers on visceral and somatic stimuli, rats, 87:382
  - peripheral, mediated by  $\kappa$ - and  $\delta$ -opioid receptors, involvement of NO, 87:388
  - somatic and visceral, tolerance, after continuous epidural infusion of morphine, rats, 87:1340
- capsaicin-induced, effects of intrathecal vs intravenous clonidine, 87:591
- catheter insertion, 87:413
- central, treatment, intrathecal R-phenylisopropyladenosine, effects of repeated administration, rats, 87:1367
- chronic benign, analgesic efficacy of morphine, effect of proglumide, 87:1117
- hyperalgesia
  - experimental thermal or capsaicin-induced, effects of intrathecal vs intravenous clonidine, normal volunteers, 87:591
  - mechanical, effects of metabotropic glutamate receptor antagonists, rats, 87:1354
- neuropathic
  - dexmedetomidine, nerve injury, and  $\alpha_2$ -adrenoceptors, 87:941
  - effects of intrathecal vs intravenous clonidine, 87:591
  - excitatory amino acid receptor antagonists, NOS inhibitor, and autotomy, 87:605
  - mechanical allodynia, anti-allodynic effects of amitriptyline, gabapentin, and lidocaine, rats, 87:1360
- noxious mechanical stimulus, simple, quantifiable, and accurate application method, 87:1446

- postherpetic neuralgia, use of percutaneous electrical nerve stimulation, 87:911
  - postoperative, management
    - abdominal surgery, intramuscular morphine at fixed rate vs epidural morphine or sufentanil and bupivacaine, 87:1346
    - acetaminophen as adjunct to morphine by PCA, 87:368
  - ambulatory surgery, use of betamethasone, 87:319
  - bilateral tubal ligation, 87:619
  - cesarean section, efficacy of intrathecal neostigmine, intrathecal morphine, and their combination, 87:341
  - cryoanalgesia, postherniorrhaphy pain, 87:896
  - fentanyl PCA, 87:890
  - intraarticular triamcinolone acetate, arthroscopic knee surgery, 87:1113
  - multimodal approach, daycase laparoscopic sterilization (letter), 87:496
  - PCEA with background infusion vs PCEA without background infusion, gastrectomy, 87:907
  - perioperative magnesium sulfate administration, 87:206
  - peripheral effect of fentanyl, 87:1121
  - peripheral nerve blocks, total knee replacement, 87:93
  - postthoracotomy, ketamine as adjunct to morphine, unintended N-of-1 study, 87:1372
  - scalp infiltration with bupivacaine, adult patients undergoing craniotomy, 87:579
  - unilateral total knee arthroplasty, 87:88
  - thermal, effects of intrathecal vs intravenous clonidine, normal volunteers, 87:591
- Pancuronium**, see Neuromuscular relaxants
- Paracetamol absorption test**, see Measurement techniques
- Parkinsonism**, see Complications
- Patient-controlled analgesia (PCA)**, see Anesthetic techniques
- Patient-controlled epidural analgesia (PCEA)**, see Anesthetic techniques
- Patient satisfaction**, see Outcome
- PCA**, see Anesthetic techniques, patient-controlled analgesia
- PCEA**, see Anesthetic techniques, patient-controlled epidural analgesia
- Pediatric anesthesia**, see Anesthesia
- Pens**, see Equipment

**Peptides**, adrenomedullin, increased plasma concentration, patients with subarachnoid hemorrhage, 87:859

**Percutaneous electrical nerve stimulation**, see Anesthetic techniques

**Peripheral nerve stimulation**, see Monitoring

# **Pharmacodynamics**

alfentanil, effect of intravenous and oral fluconazole, 87:190

cisatracurium, after short infusion, patients under propofol anesthesia, 87:1158

propofol, clearance and distribution volume increase, patients with hyperthyroidism, 87:195

# **Pharmacokinetics**

alfentanil, effect of intravenous and oral fluconazole, 87:190

cisatracurium, after short infusion, patients under propofol anesthesia, 87:1158

propofol, clearance and distribution volume increase, patients with hyperthyroidism, 87:195

# **Pharmacology**

adenosine

A<sub>1</sub>-receptor agonists, R-phenylisopropyladenosine, repeated intrathecal administration, reduced anti-allodynic effect and lack of cross-tolerance with morphine, rats, 87:1367

intracarotid, for manipulation of human cerebrovascular resistance, feasibility assessment, 87:1291

albuterol, cough during emergence from isoflurane anesthesia, 87:1170

alfentanil, reduction of MAC, potentiation by esmolol, 87:671

baclofen, withdrawal, presenting as multiorgan system failure, 87:562

benztropine, therapy, reversal of parkinsonism-induced acute upper airway obstruction, 87:975

betamethasone

effect on postoperative pain and nausea, ambulatory surgery, 87:319

topical application, to reduce sore throat, hoarseness, and coughing after tracheal intubation, 87:714

cimetidine, reduction of side effects from rapid vancomycin infusion, 87:681

clonidine

effect on recovery from myocardial stunning, dogs, 87:1009

effects on gastric intramucosal Pco<sub>2</sub>, 87:686

intrathecal clonidine and fentanyl with hyperbaric bupivacaine, effect on analgesia during cesarean section, 87:609

dexamethasone

effect on incidence of shivering after cardiac surgery, randomized, double-blind, placebo-controlled study, 87:795

preoperative, effect on immediate and delayed postoperative morbidity, children undergoing adenotonsillectomy, 87:57

dexmedetomidine, effects on neural conduction and neurotransmitter release in sympathetic ganglia, 87:503

dextromethorphan, preoperative, effect on intraoperative morphine requirements, patients undergoing laparotomy, 87:1135

dihydroergotamine, prevention of cardiovascular side effects during spinal anesthesia, 87:347

diltiazem

antinociceptive effects, visceral and somatic stimuli, rats, 87:382

pretreatment, effects on potassium release after succinylcholine administration, dogs, 87:691

diphenhydramine, reduction of side effects from rapid vancomycin infusion, 87:681

dizocilpine, vs isoflurane, relative neuroprotective effects during focal cerebral ischemia, rats, 87:72

dobutamine, vascular, metabolic, and contractile responses, influence of NO, *in situ* canine hearts, 87:994

domperidone, vs granisetron, for preventing PONV, 87:1404

donepezil, and neostigmine, effects in patient with unusual

dopamine, renal protection during CPB (letter), 87:1458

pseudocholinesterase activity, 87:1203

droperidol

granisetron-droperidol combination, prevention of PONV, children, 87:761

vs ondansetron, antiemetic effect, patients undergoing gynecologic laparoscopy (letter), 87:1213

ephedrine, pressor response during spinal anesthesia, effect of oral clonidine premedication, 87:1336

epidural drugs, new, *primum non nocere* (letter), 87:1211

esmolol, effect on reduction of MAC by alfentanil, 87:671

fibrin glue, patch, effect in *in vitro* model of postdural puncture leakage, 87:1125

glycopyrrolate, effect on incidence of hypotensive/bradycardic events during shoulder arthroscopy in sitting position under interscalene block, 87:1320

granisetron

domperidone vs, for preventing PONV, 87:1404

droperidol combination, prevention of PONV, children, 87:761

heparin

prostaglandin E<sub>1</sub> added to, increased anticoagulation during CPB, 87:985

resistance, antithrombin III treatment, and ACT values (letter), 87:1215

isoproterenol, intravenous, simulation of epidural test dose, sevoflurane- and halothane-anesthetized children, 87:549

magnesium sulfate

effect on hemodynamics and efficacy in attenuating response to endotracheal intubation, patients with coronary artery disease, 87:808

reduction of intra- and postoperative analgesic requirements, 87:206

mannitol, renal protection during CPB (letter), 87:1458

methacholine, pulmonary

responsiveness, effect of spinal anesthesia, guinea pigs, 87:874

methylprednisolone, pulmonary effects, patients undergoing CABG and early tracheal extubation, 87:27

metoclopramide, administration, followed by akathisia and refusal of surgery, 87:224

metoprolol, effect on incidence of hypotensive/bradycardic events during shoulder arthroscopy in sitting position under interscalene block, 87:1320

milrinone, effects and mechanism in fatigued diaphragm, dogs, 87:1077

nicardipine, intracarotid, manipulating human cerebrovascular resistance, 87:1291

nicotine, -induced analgesia, effects of serotonin biosynthesis inhibition, rats, 87:1109

nifedipine, -induced analgesia, effects of serotonin biosynthesis inhibition, rats, 87:1109

- nitroglycerin  
  preferential effect on large microvessels (letter), 87:747  
  pulmonary hypertension and major surgery, 87:812
- ondansetron  
  anaphylactoid reaction, pediatric neurosurgical patient, 87:779  
  droperidol vs, antiemetic effect, patients undergoing gynecologic laparoscopy (letter), 87:1213
- phenylephrine  
  hypertension induced by, failure to improve outcome after closed head trauma, rats, 87:574  
  renal protection during CPB (letter), 87:1458
- PPBP, neuroprotection from focal ischemia, dependence on treatment duration, rats, 87:1299
- progumide, effect on analgesic efficacy of morphine, humans with chronic benign pain, 87:1117
- protamine, reversal of heparin, effects on platelet aggregation and ACT after CPB, 87:781
- RS-25259, preventing PONV after hysterectomy procedures, 87:462
- triamcinolone acetonide, intraarticular, for pain control after arthroscopic knee surgery, 87:1113
- tropisetron, single-dose, preventing PONV after breast surgery, 87:931
- verapamil  
  antinociceptive effects on visceral and somatic stimuli, rats, 87:382  
  pretreatment, effects on potassium release after succinylcholine administration, dogs, 87:691
- Pharyngoscopic views**, proposed new class (letter), 87:748
- Pharynx**  
  edema, airway obstruction caused by, succinylcholine and (letter), 87:229  
  mucosal pressure, and airway sealing pressure, with LMA, comparison, anesthetized adult patients, 87:1379
- Phenylephrine**, see Pharmacology
- 4-Phenyl-1-(4-phenylbutyl)piperidine (PPBP)**, see Pharmacology
- Pheochromocytoma**, see Complications
- Plasma cholinesterase**, see Enzymes
- PONV**, see Vomiting, nausea, and, postoperative
- Popliteal fossa**, see Knee
- Position**  
  lateral decubitus, advantages of, after spinal anesthesia with hyperbaric tetracaine, 87:879  
  lateral, effects on cardiopulmonary function during laparoscopic urological surgery, 87:925  
  maternal, during induction of CSE anesthesia for cesarean delivery, 87:614  
  prone, effect on pulmonary mechanics, frame-dependence, 87:1175  
  sitting, shoulder arthroscopy under interscalene block, hypotension, preventing, use of metoprolol and glycopyrrolate, 87:1320
- Postanesthesia care unit (PACU)**  
  exposure of nurses to exhaled anesthetic gases, 87:1083  
  fast-track eligibility, after ambulatory anesthesia with propofol and desflurane, and Bispectral Index, 87:1245  
  length-of-stay, quantifying and assessing dependent factors, 87:628
- Posterior cingulate cortex**, see Brain
- Potassium**, see Ions
- Potency**, anesthetic  
  additivity vs synergism, combined spinal morphine-systemic buprenorphine, 87:579  
  inhaled compounds, anesthetic and convulsant requirements, influence of rat strain, 87:963  
  Meyer-Overton hypothesis, MACs of noble gases, 87:419  
  minimum alveolar anesthetic concentration (MAC)  
    ethanol, concentrations required to suppress learning in fear-potentiated startle paradigm, rats, 87:1398  
  isoflurane, reduction by alfentanil, potentiation by esmolol, 87:671  
  noble gases, nitrogen, and sulfur hexafluoride, 87:419  
  sevoflurane, end-tidal concentration for tracheal extubation and skin incision, children, 87:1263  
  understanding and defining (letter), 87:1453
- PPBP**, see Pharmacology
- 4-phenyl-1-(4-phenylbutyl)piperidine**, see Pharmacology
- Preemptive analgesia**, see Anesthetic techniques
- Premedication**  
  clonidine  
    effect on perioperative responses during ketamine anesthesia, 87:161  
  failure to attenuate suppression of certain lymphocyte subsets after surgery 87:1426
- lorazepam and morphine-scopolamine vs, patients undergoing CABG, 87:292  
  oral, effect on pressor response to ephedrine during spinal anesthesia, 87:1336  
  redistribution hypothermia after induction of anesthesia, 87:168  
  lorazepam, vs clonidine, patients undergoing CABG, 87:292  
  midazolam, effect on perioperative responses during ketamine anesthesia, 87:161  
  morphine, -scopolamine, vs clonidine, patients undergoing CABG, 87:292  
  scopolamine, morphine-scopolamine, vs clonidine, patients undergoing CABG, 87:292
- Preoperative period**, behavioral preparation programs, in children, comparative examination, 87:1249
- Preperitoneal fascial block**, see Anesthetic techniques
- Progumide**, see Pharmacology
- Propofol**, see Anesthetics, intravenous
- Prostaglandin E<sub>1</sub>**  
  alprostadil, increased anticoagulation during CPB, 87:985  
  intrapulmonary shunt fraction and (letter), 87:233
- Prostatectomy**, see Surgery, urological
- Protamine**, see Pharmacology
- Proteins**  
  c-fos, ketamine-induced, in posterior cingulate cortex, effect of propofol, rats, 87:1416  
  lipoproteins and serum binding variations of propofol, 87:1147
- Pruritus**, see Complications
- Pseudocholinesterase**, see Enzymes
- Pseudotongue enlargement**, see Complications
- Pulmonary aspiration**, see Complications
- Pulmonary embolism**, see Embolism
- Pulmonary fibrosis**, see Complications
- Pulmonary hypertension**, see Lung, hypertension
- Pyloric stenosis**, see Complications
- Quality of care**, patient satisfaction with anesthesia care, measuring, review of current methodology, 87:1089
- RCV**, see Measurement techniques, red cell volume
- Receptors**  
  adenosine, A<sub>1</sub>-receptor agonists, R-phenylisopropyladenosine, repeated intrathecal administration, reduced anti-allodynic effect and lack of cross-tolerance with morphine, rats, 87:1367

- $\alpha$ -adrenoceptors, stimulation, inotropic responses, modifications by propofol, rat myocardium, 87:277
- $\alpha_2$ -adrenoceptors
  - agonists, inhibition of sympathetic ganglionic transmission, mechanism, 87:503
  - agonists, safety (letter), 87:496
  - peripheral, possible role, enhancement of analgesic potency of dexmedetomidine after nerve injury, 87:941
  - stimulation, inhibition of sympathetic ganglionic transmission, mechanism, 87:503
- $\beta$ -adrenoceptors, stimulation, inotropic responses, modifications by propofol, rat myocardium, 87:277
- amino acids, excitatory, antagonists, intrathecal
  - effect on autotomy behavior, rats, 87:605
  - mechanical hyperalgesia and, rats, 87:1354
- metabotropic glutamate, antagonists, intrathecal, failure to decrease mechanical hyperalgesia, rats, 87:1354
- N-methyl-D-aspartate (NMDA)
  - antagonism, and focal cerebral ischemia, 87:72
  - antagonists, ketamine, -induced c-fos expression, in posterior cingulate cortex, effect of propofol, rats, 87:1416
  - role in relaxant effect of ketamine on tracheal smooth muscle, 87:1383
- $\kappa$ -opioid
  - partial agonistic activity of naloxone, Chinese hamster ovary cells, 87:450
  - peripheral antinociception mediated by, involvement of NO, 87:388
- $\mu$ -opioid
  - partial agonistic activity of naloxone, Chinese hamster ovary cells, 87:450
  - tramadol and thermoregulation, 87:173
- $\delta$ -opioid, peripheral antinociception mediated by, involvement of NO, 87:388
- Recovery**
  - coughing, emergence from isoflurane anesthesia, 87:1170
  - fast-track eligibility, after ambulatory anesthesia with propofol and desflurane, and Bispectral Index, 87:1245
- time
  - discharge, factors affecting, adult outpatients, 87:816
  - postoperative, impact of choice of muscle relaxant (letter), 87:499
- Red cells**, see Blood; Measurement techniques, red cell volume (RCV)
- "Red man syndrome,"** see Complications
- Reflexes**
  - hypoxic pulmonary vasoconstriction, effect of prolonged inhaled NO, rats, 87:1285
  - somatosympathetic, A $\delta$  and C, synergism between sevoflurane and intravenous fentanyl, dogs, 87:211
- Regional anesthesia and pain management**, see Anesthesia
- Remifentanyl**, see Analgesics
- Reperfusion injury**, see Heart
- Residents**, see Education
- Respiratory depression**, see Complications
- Respiratory mechanics**, see Ventilation
- Rocuronium**, see Neuromuscular relaxants
- Ropivacaine**, see Anesthetics, local
- RS-25259**, see Pharmacology
- Saline**, epidural, and combined epidural-spinal anesthesia (letter), 87:232
- Sciatic nerve block**, see Anesthetic techniques
- Scopolamine**, see Premedication
- Sedation**
  - after aortic surgery, infusion of propofol, sufentanil, or midazolam, comparison of oxygen consumption and hemodynamic stability, 87:272
  - levels, defining, pediatric patients (letter), 87:1454
- Semantics**, importance of choice of words (letter), 87:743
- Serotonin**
  - see Metabolism
  - see Neurotransmitters
- Sevoflurane**, see Anesthetics, volatile
- Sharps disposal**, see Operating room (OR)
- Shivering**, see Thermoregulation
- Shoulder surgery**, see Surgery
- Shy-Drager syndrome**, see Complications
- Sialoadenitis**, see Complications
- Sialoadenopathy**, see Complications
- Sinus surgery**, see Surgery
- Skin vasomotor reflex test**, see Measurement techniques
- Smoking**, see Complications
- Sodium**, see Ions
- Sodium fluorescein**, see Nonradioactive markers
- Somatosensory evoked potentials**, see Brain
- Somatosympathetic reflexes**, see Reflexes
- Sonoclot analysis**, see Measurement techniques
- Sore throat**, see Complications
- Spinal anesthesia**, see Anesthesia
- Spinal anesthetic solutions**, composition, variations, effects of drug addition order and preparation methods, 87:1326
- Spinal cord**
  - differential delivery of intravenous drugs, goats, 87:1450
  - dorsal horn
    - metabotropic glutamate receptors, antagonists, intrathecal, and mechanical hyperalgesia, rats, 87:1354
  - plasma and CSF lidocaine concentrations, 87:601
- Spinal cord stimulator**, see Anesthetic techniques
- Spinal fluid**, see Cerebrospinal fluid (CSF)
- Spine**, cervical vertebrae, patient with ankylosing spondylitis, successful direct laryngoscopy assisted by posture, 87:1436
- Spine surgery**, see Surgery
- Spinal motor neurons**, see Neurons
- Splenectomy**, see Surgery
- Staphylococcus aureus**, see Infection
- Statistics**
  - analysis, total labor pain, use of visual analog scale, 87:723
  - epidemiology, measuring patient satisfaction with anesthesia care, 87:1089
  - linear logistic regression, risk factors of death in traumatized children, 87:537
  - meta-analysis (letter), 87:1460
- Stellate ganglion block**, see Anesthetic techniques
- Sterilization**, see Surgery
- Stress**, surgical, influence on T cells, enhancement of early phase lymphocyte activation, 87:1431
- ST trending monitors**, see Equipment
- Subarachnoid hemorrhage**, see Complications
- Subcutaneous anesthesia**, see Anesthetic techniques
- Succinylcholine**, see Neuromuscular relaxants
- Sufentanil**
  - see Analgesics
  - see Anesthetics, intravenous
- Sulfur hexafluoride**, MACs of noble gases, 87:419

**Surgery**

## abdominal

aortic, aneurysmectomy, prophylactic albumin administration before aortic unclamping, pulmonary arterial and right ventricular responses, 87:1020

aortic, elective, patients undergoing preoperative hemodilution, hemodynamics and oxygen transport, effects of increased doses of bovine hemoglobin, 87:284

major, intramuscular morphine at fixed rate vs epidural morphine or sufentanil and bupivacaine, 87:1346

adrenalectomy, for pheochromocytoma, hemodynamic and catecholamine responses, pediatric patient, 87:1268

## aortic

abdominal, aneurysmectomy, prophylactic albumin administration before aortic unclamping, pulmonary arterial and right ventricular responses, 87:1020

abdominal, elective, patients undergoing preoperative hemodilution, hemodynamics and oxygen transport, effects of increased doses of bovine hemoglobin, 87:284

myocardial ischemia after, rates, epidural analgesia vs intravenous PCA (letter), 87:745

sedation after, infusion of propofol, sufentanil, or midazolam, comparison of oxygen consumption and hemodynamic stability, 87:272

## cardiac

evaluation of platelet function by Sonoclot analysis compared with other hemodynamic variables, 87:1228

postoperative analgesia, nurse-administered subcutaneous morphine as alternative to intravenous PCA morphine, 87:11

postoperative shivering, effect of dexamethasone, 87:795

repair of congenital heart defects, thyroid axis, effects of CPB and deep hypothermic circulatory arrest, children, 87:543

repair of congenital heart disease, effects of inhaled NO on pulmonary hypertension, infants and children, 87:46

## cardiopulmonary bypass (CPB)

effects on thyroid axis during and after repair of congenital heart defects, children, 87:543

increased anticoagulation by prostaglandin E<sub>1</sub>, 87:985

IL-6 response after, effect of ketamine, 87:266

interpretation of nonpulsatile arterial pressure-flow relations during (letter), 87:743

postoperative bleeding and transfusions, effects of tranexamic acid vs aprotinin, 87:258

postoperative platelet aggregation and ACT, effect of protamine reversal of heparin, 87:781

renal protection during, use of phenylephrine, dopamine, and mannitol (letter), 87:1458

carotid endarterectomy, mandibular nerve block in addition to cervical plexus block, 87:1034

## cesarean section

analgesia during, effect of intrathecal clonidine and fentanyl with hyperbaric bupivacaine, 87:609

CSE anesthesia, effects of maternal position during induction, 87:614 for dystocia, incidence, and epidural labor analgesia, 87:119

intrathecal hypobaric vs hyperbaric bupivacaine with morphine, 87:336

postoperative analgesia, combination of diclofenac and intrathecal morphine (letter), 87:1453

postoperative analgesia, efficacy of intrathecal neostigmine, intrathecal morphine, and their combination, 87:341

spinal anesthesia, hyperbaric 0.75% vs hyperbaric 1% bupivacaine, 87:1099

cholecystectomy, laparoscopic, respiratory mechanics, effects of abdominal wall lift, 87:1393

coronary artery bypass grafting (CABG) clonidine vs conventional

preanesthetic medication, 87:292

early tracheal extubation, patients undergoing, pulmonary effects of methylprednisolone, 87:27

craniotomy, hemodynamics and postoperative pain, influence of scalp infiltration with bupivacaine, adult patients, 87:579

endoscopy, upper, gastric contents, children, 87:757

facial laser resurfacing, outpatient, TIVA technique, 87:827

gastrectomy, postoperative pain, PCEA with background infusion vs PCEA without background infusion, 87:907

## gynecological

ambulatory, perioperative

hypothermia, effect of warming intravenous fluids, 87:37

hysterectomy, PONV, preventing, efficacy of RS-25259, 87:462

laparoscopic, antiemetic effect of ondansetron vs droperidol (letter), 87:1213

laparoscopic, excretion of serotonin metabolite, 87:1104

PONV, preventing, granisetron vs domperidone, 87:1404

hallux valgus repair, sciatic and femoral nerve blockade, comparison of ropivacaine, bupivacaine, and mepivacaine, 87:597

heart transplantation, patient with latex allergy, perioperative management, 87:304

## hernia repair

laparoscopic, extraperitoneal, analgesic effect of bupivacaine, 87:377

postoperative analgesia, preemptive caudal bupivacaine and morphine, children, 87:52

postoperative pain, effect of cryoanalgesia, 87:896

## hip

continuous spinal anesthesia, Shy-Drager syndrome, 87:396

total hip arthroplasty, delayed neurological deficit after, 87:1209

kidney transplantation, living donor, patients undergoing, renocortical tissue oxygen pressure measurements, 87:474

## knee

arthroscopic, postoperative pain control, use of intraarticular triamcinolone acetonide, 87:1113

total knee arthroplasty, unilateral, postoperative pain and knee rehabilitation, effects of intravenous PCA with morphine, continuous epidural analgesia, and continuous three-in-one-block, 87:88

total knee arthroplasty, without pneumatic tourniquet use, incidence of large venous emboli, 87:439

total knee replacement, postoperative analgesia, improvement, use of peripheral nerve blocks, 87:93



- laryngeal microsurgery, high-frequency jet ventilation, efficiency of CO<sub>2</sub> elimination, 87:180
- liver transplantation  
graft nonfunction, detection, assessment of indocyanine green kinetics, 87:34  
orthotopic, torsade de pointes during, 87:300
- mastectomy, PONV, prevention, single-dose tropisetron, 87:931
- middle ear, with propofol or isoflurane anesthesia, autoregulation of inner ear blood flow during controlled hypotension, 87:1002
- neurosurgery, controlled continuous spinal drainage, use of flow-regulated pump, 87:1306
- open heart, complications of CO<sub>2</sub> flooding surgical field (letter), 87:978
- radial artery cannulation, topical lidocaine-prilocaine cream (EMLA) vs local infiltration anesthesia, 87:403
- shoulder, arthroscopic, in sitting position under interscalene block, hypotensive/bradycardic events, prevention, use of metoprolol and glycopyrrolate, 87:1320
- sinus, endoscopic, postoperative "man-in-the-barrel" syndrome, 87:1196
- spine  
prone position, effect on pulmonary mechanics, frame-dependence, 87:1175  
spinal fusion, PCA morphine, dose-response of ketorolac as adjunct, 87:98
- splenectomy, elective, anemic Jehovah's Witness patient with cirrhosis, 87:529
- sterilization, laparoscopic, daycase, postoperative pain control, multimodal approach (letter), 87:496
- thoracoscopic  
arterial blood gases, substantial changes missed by conventional intermittent laboratory blood gas analyses, 87:647  
video-assisted, interruption of PDA, effects on cardiopulmonary function, newborn pigs, 87:1037
- thoracotomy  
intercostal nerve blockade, effect of adding clonidine to anesthetic solution, 87:107
- postoperative analgesia, ketamine as adjunct to morphine, unintended N-of-1 study, 87:1372
- systemic and pulmonary hemodynamics and arterial oxygenation, effects of desflurane vs isoflurane, patients undergoing one-lung ventilation, 87:800
- tonsillectomy, pediatric patients  
adenotonsillectomy, immediate and delayed postoperative morbidity, effect of preoperative dexamethasone, 87:57  
PONV, prevention, granisetron-droperidol combination, 87:761
- tracheostomy  
laryngeal trauma, anesthetic implications, 87:1283  
percutaneous, dilational, Ciaglia vs Rapitrach method, 87:556
- tubal ligation, bilateral, effective analgesia after, 87:619
- urological  
laparoscopic, cardiopulmonary function, effect of lateral position, 87:925
- prostatectomy, radical, overnight hospitalization after, impact of two clinical pathways, 87:62
- transurethral resection of prostate, spinal vs general anesthesia, ratio of T helper 1 to T helper 2 cell subsets, 87:1421
- vascular  
abdominal aortic aneurysmectomy, prophylactic albumin administration before aortic unclamping, 87:1020
- pulmonary hypertension and major surgery, 87:812
- Sweating**, see Thermoregulation
- Sympathetic ganglia**, transmission,  $\alpha_2$ -adrenergic inhibition, mechanism, 87:503
- Sympathetic nervous system**  
catecholamines  
epinephrine, postoperative plasma concentrations, effect of premedication with clonidine, 87:1426  
epinephrine, tocolytic effect, gravid rat uterus, 87:864  
norepinephrine, for cardiopulmonary resuscitation during epidural block (letter), 87:1216  
norepinephrine, postoperative plasma concentrations, effect of premedication with clonidine, 87:1426
- norepinephrine, tocolytic effect, gravid rat uterus, 87:864
- norepinephrine, tramadol and thermoregulation, 87:173
- plasma, measurement, during amino acid-induced thermogenesis, determination of stress response, 87:637
- plasma, measurement, effects of clonidine on recovery from myocardial stunning, dogs, 87:1009
- responses to laparoscopic adrenalectomy for pheochromocytoma, pediatric patient, 87:1268
- Syringes**, see Equipment
- Tachycardia**, see Heart
- Technology assessment**, overview, 87:1271
- Temperature**, hypothermia  
deep, preservation of thyroid function, children undergoing repair of congenital heart defects, 87:543  
during anesthesia, prevention, amino acid-induced thermogenesis  
postoperative nitrogen excretion, 87:641  
stress response, 87:637  
effects of tramadol on thermoregulatory responses, 87:173  
perioperative, reduced by warming intravenous fluids, women undergoing ambulatory gynecological surgery, 87:37  
redistribution, after induction of anesthesia, and clonidine premedication, 87:168
- Tetanic fade**, see Measurement techniques
- Tetracaine**, see Anesthetics, local
- Thermogenesis**, amino acid-induced to prevent hypothermia during anesthesia, and stress response, 87:637  
under anesthesia, postoperative nitrogen excretion, 87:641
- Thermoregulation**  
shivering  
after cardiac surgery, effect of dexamethasone, 87:795  
threshold, during epidural lidocaine anesthesia, effect of epidural fentanyl, 87:587  
threshold, effect of hypoxemia, rabbits anesthetized with 0.2 MAC isoflurane, 87:1408  
threshold, effect of tramadol, 87:173  
sweating, threshold, effect of tramadol, 87:173  
vasoconstriction, threshold, effect of tramadol, 87:173

- Thiopental**, see Anesthetics, intravenous
- Thoracic paravertebral blockade**, see Anesthetic techniques
- Thoracotomy**, see Surgery
- Thromboelastography**, see Measurement techniques
- Thromboembolism**, see Embolism
- Thyroid**, function, effects of CPB and deep hypothermic circulatory arrest, during repair of congenital heart defects, children, 87:543
- Tonsillectomy**, see Surgery
- Topical anesthesia**, see Anesthetic techniques
- Torsade de pointes**, see Complications
- Total intravenous anesthesia (TIVA)**, see Anesthetic techniques
- Total laryngeal bypass device**, see Equipment
- Toxicity**  
lidocaine, 87:900  
sevoflurane, 87:468
- Trachea**  
diameter, predicting size of double-lumen endobronchial tube, 87:158  
smooth muscle, relaxant effect of ketamine, role of NMDA receptor, 87:1383
- Tracheostomy**, see Surgery
- Train-of-four ratios**, see Measurement techniques
- Tramadol**, see Analgesics
- Tranexamic acid**, see Antifibrinolytics
- Transcranial Doppler ultrasonography**, see Measurement techniques
- Transcutaneous electrical nerve stimulation (TENS)**, see Anesthetic techniques
- Transdermal nicotine system**, see Complications
- Transesophageal echocardiography (TEE)**, see Monitoring
- Transfusion**, see Blood
- Transurethral resection of prostate**, see Surgery, urological
- Trauma**, see Complications
- Triamcinolone acetonide**, see Pharmacology
- Triglycerides**, serum protein binding of propofol, 87:1147
- Tropisetron**, see Pharmacology
- Tubal ligation**, see Surgery
- d-Tubocurarine**, see Neuromuscular relaxants
- Ulnar nerve**, see Nerves
- Ultrasonography**, see Measurement techniques, transcranial Doppler ultrasonography
- Univent tubes**, see Equipment
- Urethral profile**, see Measurement techniques
- Urine**  
nitrogen excretion, postoperative, after amino-acid-induced thermogenesis under anesthesia, 87:641  
serotonin metabolite excretion, gynecologic laparoscopic surgery, 87:1104
- Urodynamics**, effects of intravenous opioids and ketoprofen, humans, 87:456
- Urological surgery**, see Surgery
- Uterus**, activity, tocolytics, 87:864
- Vancomycin**, see Antibiotics
- Vasoconstriction**  
see Lung  
see Thermoregulation
- Vasoconstrictors**  
5-hydroxytryptamine, tramadol and thermoregulation, 87:173  
prevention of cardiovascular side effects during spinal anesthesia, 87:347
- Vasodilation**, see Lung
- Vecuronium**, see Neuromuscular relaxants
- Veins**, jugular, internal, cannulation, use of TEE probe imaging to guide, 87:1032
- Ventilation**  
hyperventilation brain injury, 87:850  
propofol and, for treatment of increased intracranial pressure, rabbits, 87:564  
spinal motor neuron excitability during isoflurane anesthesia, 87:407  
effect on cerebral venous oxygen saturation, patients with traumatic hypoventilation, and spinal motor neuron excitability during isoflurane anesthesia, 87:407
- jet**  
high-frequency, laryngeal microsurgery, efficiency of CO<sub>2</sub> elimination, 87:180  
in upper airway obstruction, new jetting device, description and model lung testing, 87:915  
transtracheal, mask airway and rescue ventilation, 87:661  
respiratory mechanics  
during general anesthesia, effect of body mass, 87:654  
during laparoscopic cholecystectomy, effects of abdominal wall lift, 87:1393  
pulmonary effects of spinal anesthesia, 87:874  
spontaneous, respiratory effects of desflurane anesthesia, infants and children, 87:1052  
volume vs pressure cycle, infant lung model, 87:766
- Verapamil**, see Pharmacology
- Video**, see Education, patients
- Video-assisted surgery**, see Surgery, thoracoscopic
- Visual analog scale**, see Measurement techniques
- Vomiting**, nausea and, postoperative (PONV)  
ambulatory surgery  
drinking fluids after, 87:306  
effect of betamethasone, 87:319  
effect of preoperative dexamethasone, children, 87:57  
emesis, and N<sub>2</sub>O, correction (letter), 87:1215  
gynecologic laparoscopy and serotonin excretion, 87:1104  
ondansetron vs droperidol, antiemetic effect, patients undergoing gynecologic laparoscopy (letter), 87:1213  
preventing  
after breast surgery, use of single-dose tropisetron, 87:931  
after hysterectomy procedures, efficacy of RS-25259, 87:462  
granisetron-droperidol combination, children, 87:761  
granisetron vs domperidone, 87:1404
- Xenon**, see Anesthetics, gases

